

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

PRACTICE PARAMETERS FOR THE ASSESSMENT AND TREATMENT OF CHILDREN AND ADOLESCENTS WHO ARE SEXUALLY ABUSIVE OF OTHERS

These parameters were developed by Jon A. Shaw, M.D., principal author, and the Work Group on Quality Issues: William Bernet, M.D., Chair, John Dunne, M.D., former Chair; Maureen Adair, M.D., Valerie Arnold, M.D., Joseph Beitchman, M.D., R. Scott Benson, M.D., Oscar Bukstein, M.D., Joan Kinlan, M.D., Jon McClellan, M.D., and David Rue, M.D. AACAP Staff: Elizabeth Sloan, L.P.C., and Kristin D. Kroeger. The authors thank William D. Erickson, M.D., John A Hunter, Ph.D., Toni Cavanaugh Johnson, Ph.D., and Barry M. Maletzky, M.D., for their thoughtful review. These parameters were made available to the entire Academy membership for review in September, 1998 and were approved by the Academy Council on June 27, 1999. They are available to AACAP members on the World Wide Web, (WWW.aacap.org.)

Reprint requests to AACAP, Communications Department, 3615 Wisconsin Avenue, N.W., Washington, D.C. 20016.

(c)1999 by the American Academy of Child and Adolescent Psychiatry.

ABSTRACT

The assessment and treatment of children and adolescents with sexually abusive behavior requires an understanding of normal sexual development. A multiplicity of biological and psychosocial factors determines the child's sexual development, gender role, sexual orientation, patterns of sexual arousal, sexual cognitions, sexual socialization and the integration of sexual and aggressive patterns of behavior. The individual's sexuality evolves in concert and as a result of interaction with family, ethnic, social and cultural influences. These parameters summarize what we know about the epidemiology and phenomenology of sexually abusive youth and provide guidelines for the assessment and the selection of treatment interventions for these youths. Essential considerations in the assessment and treatment of sexually abusive youth, as well as the different categories of sexually abusive youth that should be recognized and which influence treatment decisions are presented. The spectrum of currently available psychosocial and biological treatments will be summarized. **Key Words:** juvenile, sex, sexual, abuse, abuser, offender, offenses.

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

INTRODUCTION

There is evidence of a significant increase in the reports of juvenile sexual aggression and sexual abuse. Sexual assault is one of the fastest growing violent crimes in the United States. Approximately one out of three women and one out of every seven men will be sexually victimized before 18 years of age.

Studies of adult sex offenders have demonstrated that the majority self-report the onset of sexual offending behavior before 18 years of age. Approximately 20% of all rapes and 30-50% of child molestations are carried out by youth under 18 years of age. Studies of adolescent sex offenders have shown that the majority commit their first sexual offense before 15 years of age and not infrequently before 12 years of age. There are increasing reports of preadolescent sexual abusers.

The recognition of a group of children and adolescents who sexually abuse others requires that we begin to develop empirically based methods of assessment and intervention strategies so that the sexually abusive youth of today do not become the adult sex offenders of tomorrow.

EXECUTIVE SUMMARY

THE EVALUATION PROCESS

The evaluation and treatment of children and adolescents with sexually abusive behavior require an understanding of the biological and psychosocial factors determining the child's sexual development, gender role, sexual orientation, patterns of sexual arousal, sexual cognitions, sexual socialization and the integration of sexual and aggressive patterns of behavior. The individual's sexuality evolves in concert and as a result of interaction with family, ethnic, social and cultural influences.

The clinical assessment of juvenile sexual abusers requires the same comprehensive evaluation as other children and adolescents. Important sources of information include medical and psychological reports, offense reports, victim statements, protective services reports, and probation reports. The collateral information should be obtained before the individual interview; otherwise one is left relatively unprepared before the offender's normal proclivity to minimize and deny.

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

Forensic Considerations

It is essential that the clinician define his role in the assessment of the sexual abuser. While many of the issues are relevant to a forensic evaluation, these parameters are designed to provide guidelines for the clinical evaluation of the sexual abuser. It is generally preferable to conduct the clinical evaluation after adjudication. The focus of the clinical interview is on assessing amenability to treatment, required levels of care, treatment goals and objectives and the risk of reoffending. The juvenile sex abuser is advised of reporting laws and the limits of professional confidentiality. An informed consent signed by the juvenile and his parent/guardian should be obtained prior to the clinical interview. It is important to educate and clarify for the individual and his family what is going to happen and when. The role of protective services and the juvenile justice system should be explained when relevant. Consent forms should be developed to cover the use of controversial assessment and treatment procedures such as phallometric assessment, aversive conditioning, and medications that are not accepted as standard of practice.

THE CLINICAL INTERVIEW

The clinical interview is the cornerstone of the evaluation of juvenile sexual abusers. It is necessary to establish the nature of the sexually abusive behavior. Since laws have frequently been transgressed the offender is often less than forthcoming. Issues of shame, guilt and fear of punishment impede disclosure. The clinician is advised to adopt a nonjudgmental stance and to relate to the juvenile offender in a matter-of-fact exploratory manner. The clinician clarifies the meaning of sexual jargon and avoids its use. The interview is initiated with a nonthreatening line of questioning thereby minimizing the initial defensiveness. The interviewer develops lines of questioning that will allow him/her to know more about the offender, his family, school and his current life situation. Confront minimization, denial and the apparent omissions of important information. There is little value in getting angry and accusatory. It is more useful to be patient, persistent and not easily dissuaded.

The Assessment of the Sexually Abusive Behavior

In the evaluation the clinician develops certain lines of questioning regarding the sexual abuser and the sexual abusive incident.

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

Insert Table 2 Here

The Sexual-Aggressive History. The clinician obtains the juvenile's sexual history and assesses his sexual knowledge and education, sexual development, and sexual experiences. Inquiries are directed to learn what the juvenile knows about gender differences, sexual intercourse, and his preferred patterns of sexual behaviors. Specific questions may be asked regarding the juvenile's understanding and knowledge of normal sexual activities, i.e., kissing, dating, petting, masturbation, whether he has been sexually active and engaged in intercourse, homoerotic experiences, etc. Has he been exposed to inappropriate and explicit sexuality? The clinician attempts to delineate the established pattern and spectrum of previously committed sexual aggressive acts; the victim profile; the internal and external triggers that initiate the sexual abuse cycle; the role of aggression and sadism in the sexual offense; the need to dominate, control, and humiliate the victim; the erotization of the aggression; the history of sexual victimization, physical abuse, and emotional neglect; and the history of prior non-sexual delinquent behavior. It is necessary to discriminate between compulsive sexual behaviors and paraphilic compulsive sexual behaviors. Is there a history of arrests, convictions, incarcerations, use of weapons or cruelty to animals?

Developmental and Psychosocial History. Other areas of the assessment process are those associated with a comprehensive developmental history, i.e., the nature of the pregnancy, perinatal history, developmental milestones, family relationships, early identificatory models, capacity for relationships, peer relationships and social skills. The family assessment provides an opportunity to understand the early developmental and environmental context within which the sexual abuser developed. Information is obtained regarding the parents' personal and psychological history, their use of authority and discipline, the role of coercive sexuality in the family. How is affection, tenderness, competition, aggression, love, sexuality and lust expressed in the family? How supportive and available is the family as a treatment resource?

Medical and Psychiatric History. It is important to obtain a comprehensive medical and psychiatric history with specific attention to psychopathology, substance abuse and psychiatric co-morbidity.

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

School History. A specific area of concern is the evaluation of intellectual capacities, and academic performance. Fifty to 80% of juvenile sexual abusers have learning problems, repeated a grade in school, and/or have been in classes for the learning disabled.

Mental Status Examination. A comprehensive mental status examination is carried out to assess the presence of psychopathology, personality disturbances, organicity, and substance abuse and to acquire an understanding of adaptive, coping, and defensive strategies. Suicidal content and risk should be assessed specifically. A careful assessment of the spectrum of suicidal behavior is undertaken to establish the degree to which suicidal ideation, history of suicidal behaviors, threats or plans are present. Apprehension by judicial authority and the associated shame of exposure, embarrassment, stigmatization, fear of punishment and incarceration are risk factors for suicidal behavior.

Psychological Tests

There are no specific empirical measures or psychometric tests that can identify, diagnose or classify sexual abusers. Although psychological testing may be used adjunctively to understand the personality traits, sexual behaviors, and intellectual capacities of these youngsters.

Phallometric Assessment

Some authors have recommended the use of phallometric testing, the measuring of penile erection in response to different stimuli as a way to determine sexual preferences. This technique is usually reserved for the most severe and older juvenile sexual abusers. This procedure has generally been used with caution because of the lack of empirical studies, problems in obtaining informed consent, and a reluctance to expose children and adolescents to further sexual stimulation through the portrayal of deviant sexual activities.

Disposition

At the end of the assessment process, the clinician should be prepared to address the following issues and to provide guidance to other professionals, the juvenile court and community agencies.

- The risk of repeating the sexually aggressive behavior
- The treatment needs of the individual and his family

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

- The appropriateness of removing the sexual abuser from his family
- The appropriate treatment program for the abuser, e.g. a community outpatient treatment program or a more restrictive environment, such as a detention center, residential program, or inpatient unit.

In the final stages of the evaluation, it is useful to discuss the possible treatment alternatives with the patient and appropriate family members. Explain to the family members what their participation will be in the treatment program.

TREATMENT

The spectrum of emotional, behavioral and developmental problems presented by these young people requires an integrated multi-model treatment program which is tailored to the individual's clinical presentation, and social and family support system. The treatment of juvenile sexual abusers has generally focused on several objectives, which are integral to a successful intervention:

- Confronting the sex abuser's denial
- Decreasing deviant sexual arousal
- Facilitating the development of non-deviant sexual interests
- Promoting victim empathy
- Enhancing social and interpersonal skills
- Assisting with values clarification
- Clarifying cognitive distortions
- Teaching the juvenile to recognize the internal and external antecedents of sexual offending behavior with appropriate intervention strategies

The predominant treatment approaches include cognitive-behavioral, and psychosocial therapies and psychopharmacological interventions. A group setting is the preferred format in treatment programs for sexual abusers and is usually the conduit through which cognitive-behavioral modalities (such as psycho-educational, behavioral, and relapse prevention programs) are conducted.

Cognitive-Behavioral Interventions

Psychoeducational modules provide information about sexuality, sexual deviancy, cognitive distortions, interpersonal and social behaviors, and strategies for coping with aggressive and sexual impulses. Specific educational modules may include: victim

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

awareness/empathy; cognitive restructuring; anger management; assertiveness training; social skills training; sexual education; stress reduction and relaxation management; and autobiographical awareness.

Specific behavioral techniques that have been used to diminish deviant sexual arousal include: covert sensitization; assisted covert sensitization; imaginal desensitization; olfactory conditioning; satiation techniques; and sexual arousal reconditioning.

Relapse prevention assumes that sexual offenses are the product of contextual triggers and an array of emotional and cognitive precursors. In this intervention, the sexual abuser becomes aware of each phase of his sexual assault cycle and its unique characteristics, so that he will become knowledgeable about the triggers that initiate the cycle. The goals of relapse prevention are to empower the offender to manage his own sexual life through a cognitive understanding of the antecedents of his sexual offending behavior and the development of coping strategies with which to interrupt the sexual offending cycle.

Psychosocial Therapies

Psychosocial therapies include traditional individual approaches, family therapy, group therapies, and the use of the therapeutic community.

Group therapy with juvenile sex offenders provides a context in which the sexual abuser is unable to easily minimize, deny, or rationalize his sexual behaviors. Peer group therapy, as the medium for therapeutic interventions, is used in a number of different ways depending on the setting, group membership, the severity of the sexual offenses, group goals and objectives, whether the groups are open or closed, and the length of the group experience. Therapeutic community groups are often used in hospital or residential treatment settings as a vehicle for milieu administrative decision-making and for the monitoring of a behavioral management system.

Family therapy may be most useful in those instances where there is incest, especially when the sex offender remains in the family or will rejoin the nuclear family after treatment. Family therapy facilitates the learning of new ways of communicating and building a support system which will help interrupt the abuse cycle and ultimately be supportive to the offender's capacity for regulating and modulating sexual aggressive behavior. The parents should be seen for counseling or be placed in a concurrent structured parent group with an emphasis on educational modules where they can become familiar with sexual abusive behavior, risk and

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

protective factors, characteristics of sexual abusers, treatment strategies, and most importantly focus on styles of interaction and management of their children's sexual behavior.

Individual therapy is usually employed in conjunction with other treatment approaches and probably should never be relied on as the only treatment modality. However, individual therapy may be the treatment of choice for the younger sexually-reactive abused child who has become sexually abusive. This is particularly true for children who manifest high levels of intra psychic conflict, emotional distress, confusion, and defensiveness around their own sexual victimization.

Psychopharmacological Interventions

Selective serotonin reuptake inhibitors (SSRIs) have been shown to have an impact on sexual drive, arousal and sexual preoccupations. Fluoxetine has been the agent most studied and there are a number of reports indicating that its use is associated with a reduction in paraphiliac behavior and non-paraphiliac sexual obsessions.

The antiandrogen drugs are reserved for the most severe sexual abusers and are generally discouraged for use in adolescents younger than 17 years of age. Antiandrogen medications should never be used as an exclusive treatment for paraphiliac and aggressive sexual behaviors.

LITERATURE REVIEW

A computer literature search based on Medline and Psychological Abstracts using key words such as child, adolescent, juvenile, sex, sexual, abuse, abuser, offenses and offender was obtained. References included major review articles, book chapters and monographs as well as journals with a specific focus on sexual abusive behaviors. Additionally the authors and consultants contributed from their own cumulative clinical and professional experiences.

DEFINITIONS

The more commonly used terms in the sexual abuse literature and their definitions are noted in Table 1.

Insert Table 1 Here

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

Sexually abusive behavior occurs without consent; without equality; or as a result of coercion (National Task Force on Juvenile Sexual Offending, 1993). In this context consent is defined as including all of the following: a) understanding what is proposed; b) knowledge of societal standards for what is proposed; c) awareness of potential consequences and alternatives; d) assumption that agreement or disagreement will be respected equally; e) voluntary decision; and, f) mental competence. Equality is defined as “two participants operating with the same level of power in a relationship, neither being controlled or coerced by the other.” Coercion is defined as “exploitation of authority, use of bribes, threats of force, or intimidation to gain cooperation or compliance.”

NORMAL SEXUAL DEVELOPMENT

The developing child learns about sexuality, internalizes sexual values and enacts various sexual roles as a result of exposure to familial, societal and cultural experiences. Traditionally, society has placed restraining influences on childhood sexuality. There has been a tendency to avoid sexual stimulation, to inhibit sexual impulses, to prohibit erotic play, and reduce or forbid sexual self-stimulation (Rosenfeld and Wasserman, 1993). A majority of children, however, will engage in some manifest sexual behaviors and sexual activities with others before thirteen years of age (Friedrich et al., 1991; Johnson, 1999; Araj, 1997). The sexual life of children begins to configure shortly after birth and becomes patterned upon the bases of early sensitizing experiences. In the first year of life, most children discover the pleasure of genital self-stimulation. By three to four years of age, children may begin to engage in sexual play with peers. Penile erections, thigh rubbing in female preschoolers, sexual exploration games, touching and rubbing of one's genitals, exhibitionism, voyeurism, use of dirty language and flirtatious behaviors have been described in normal children 2-6 years of age (Friedrich et al., 1991; Araj, 1997). Sexual interests during the middle childhood years wax and wane with the degree of sexual stimulation and sexually sensitizing experiences. Kissing and holding hands may occur. Sexual play between children such as “playing doctor” is normal and only becomes a concern when coercion occurs and there is an absence of mutual consent. In the process of growing up, children are invariably sexually stimulated and sometimes sexually aroused. They continually seek sexual information and greater understanding about the nature of sexual life. Through play and sexual exploration with others, as well as through gender role

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

enactments, the child begins to assimilate the elements of sexual life and establish patterns of sexual excitement and pathways to sexual gratification. Most sexual play is between children who have an ongoing mutually enjoyable play relationship and/or school friendship. The child's interest in sex and sexuality is balanced by curiosity about other aspects of his or her life. Sexual behaviors of children vary greatly and are influenced by fortuitous and opportunistic experiences, the degree of sexual stimulation, the child's sexual interest and curiosity, as well as the child's previous sexual experiences (Johnson, 1999). Normative sexual play is usually spontaneous, and includes pleasure, joy, laughter, embarrassment, and varying levels of inhibition and disinhibition (Araji, 1997). Masturbatory behavior becomes more frequent in pre adolescence. Masturbation is considered excessive when the child's masturbatory practice leads to pain or bruising or occurs in public.

Money (1986) has suggested that a relatively stable and preferred pattern of sexual gratification and erotic imagery can occur as early as eight years of age. He introduced the concept of the "lovemap." The lovemap is conceptualized as a developmental representation in the mind. It is a schema depicting the preferred sexual object and the preferred sexual-erotic behavior which evolves out of the developmental experience with others.

While early development provides the crucible for the development of sexual values, it is apparent that sexual development is greatly facilitated during the middle school years when children are increasingly exposed to the popular culture (Postman, 1994). In contemporary society children and adults have virtually equal access to sexually explicit information vis-a-vis videotapes, the Internet, television programming, and pornographic magazines. The age of "electronic information" has resulted in the popular culture in many instances superceding the family as the source of information about what is acceptable sexual behavior (Postman, 1994).

EPIDEMIOLOGY

Acts of sexual aggression have become increasingly commonplace. There has been an increase in both violent crimes committed by juveniles (OJJDP, 1994) and in the reports of sexual aggression and sexual abuse (Hampton, 1995). The National Committee to Prevent Child Abuse (1998) estimates that of the one million cases of confirmed child maltreatment in 1997 that approximately 8% were the victims of sexual abuse. Sexual assault is one of the fastest growing violent crimes in the United States and accounts for 7% of all violent crimes (Hampton, 1995).

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

While adolescents (15-18 years) make up only 6% of the population of the United States, they commit 25% of the "index crimes," such as arson, homicide, manslaughter, robbery, aggravated assault, burglary, larceny, and forcible rape (Siegel and Senna, 1988). From 1983 to 1992 there was a reported 8-10% increase in the proportion of adolescents involved in some type of serious violent offense (Elliott, 1994a). Forcible rape arrests by juveniles increased 20% percent between 1983-1992 (OJJDP, 1994). Violent behavior is highly correlated with being male at a specific phase of life. The highest risk for the initiation of serious violent behavior occurs between 15 and 16 years (Elliott, 1994a). In Elliott's (1994a, 1994b) longitudinal study of a national probability sample of 1,725 youth (11-17 years) he found that the prevalence of serious violent offenses such as rape that involved some injury or use of a weapon peaked at 17 years of age for males and 15-16 years for females. Three gender differences were apparent:

- Violent behavior occurs earlier in females
- The subsequent decline in violent behavior is steeper for females
- The gender differences becomes greater over time.

The increase in the reports of youthful sexual aggression has been well documented (Elliott, 1994a, 1994b, 1994c; OJJDP, 1994a). Ageton (1983) concluded from a probability sample of 863 male adolescents 13 to 19 years of age, that the rate of sexual assault per 100,000 adolescent males ranged from 5,000 to 16,000. A survey of high school students revealed that 1 out of 5 had been involved in forcing sex upon another, and that 60% of the boys found it acceptable in one or more situations for a boy to force sex on a girl (Davis et al., 1993). Ryan et al. (1996) in her national survey of 1,600 sexually abusive youth reported that they came from all racial, economic, ethnic, religious backgrounds and generally mirrored the population. They ranged in age from 5 to 21 years with a modal age of 14 years and were predominantly male. While the majority of juvenile sexual abusers are adolescents there is increasing concern about the sexually aggressive and exploitative behaviors in prepubescent and latency age children (Araji, 1997; Gil and Johnson, 1993; Johnson, 1988).

THE SPECTRUM OF SEXUAL OFFENSES

The spectrum of sexually inappropriate behaviors ranges from various forms of sexual harassment and non-contact sexual behaviors such as obscene phone calls, exhibitionism, and voyeurism to varying degrees of sexual aggression involving direct sexual contact including

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

frottage, fondling, digital and penile penetration, fellatio, sodomy, and other aggressive sexual acts.

A form of sexual aggression which is becoming more visible is child-on-child sexual harassment, which is unwanted sexual attention being directed to other children and adolescents. Child-on-child sexual harassment has become increasingly evident in our schools. Straus (1988) found that approximately 50% of high school junior and senior girls reported incidents of sexual harassment. Ponton (1996) reports that sexual harassment is found throughout the school environment and that student-to-student harassment is more common than adult-to-student harassment. It is known that child victims of harassment experience an array of emotional and behavioral effects (Ponton, 1996).

The spectrum of sexual offending behavior reported in the literature varies depending on the sample studied. For example, children and adolescents sampled in detention centers, residential treatment programs and outpatient clinics report different spectra of sexually offensive behavior.

Ryan et al. (1996) found in her survey of sexually abusive youth from a diversity of outpatient and residential programs that they had participated in a wide range of sexual offenses. Seventy percent of the sexual offenses involved penetration and/or oral genital behavior: vaginal or anal penetration without oral-genital contact, 35%; oral-genital contact, 14.7; penetration and oral-genital contact, 18%.

Studies of outpatient populations of juvenile sexual abusers indicate that the most common sexual offenses are fondling or "indecent liberties," 40-60%; rape and/or sodomy, 20-40%; and non-contact sexual offenses, 5-10% (Fehrenbach et al., 1986; Smith and Monastersky, 1986). The average juvenile sex offender under 18 years of age has committed 8 to 9 sexual offenses and averaged 4 to 7 victims (Abel et al., 1986 ; Shaw et al., 1993).

While adolescents usually employ coercion in the process of committing sexual offenses they are less likely to physically harm their victims when compared to adult sex offenders. The coercion is usually expressed as bribery, intimidation, threats of harm or violent injury, physical force and rarely the use of a weapon (Fehrenbach et al., 1986, Knight and Prentky, 1993). Nevertheless, it is well known that victims report higher levels of coercion and force than is self-reported by offenders (Davis and Leitenberg, 1987). Fehrenbach et al. (1986) found that 22% of the offenders continued their sexual aggressive acts even when the victims expressed "hurt or fear."

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

VICTIM CHARACTERISTICS

The great majority of the victims of male sex offenders are females (Longo, 1982; Fehrenbach et al., 1986; Davis and Leitenberg, 1987; Ryan et al., 1996). Adolescent sex offenders commit most of the sexual assaults against boys (Hunter and Becker, 1999). Ninety percent of sexual abuse victims are between 3 and 16 years of age. The majority of the victims of juvenile sex offenders are younger than 9 years of age and approximately 25-40% are under 6 years of age (Abel and Osborn, 1992; Fehrenbach et al., 1986; Ryan et al., 1996). Boys, when victimized, tend to be younger than their female counterparts (Hunter, 1991).

Distinctions have been made between those juveniles who commit incest and those who select victims outside of the family. Approximately 40 percent of the victims are blood related. Worling (1995) compared adolescent sex offenders who victimized siblings in contrast to those who selected victims outside of family. He found that those who committed incest came from families with significantly more marital discord, parental rejection, physical discipline, negative family atmosphere and were generally more dissatisfied with their family relationships. These offenders were more likely to have been sexually victimized.

It has generally been believed that incest offenders are exclusively incestuous and thus more treatable (Becker et al., 1993; Becker, 1994). Abel et al. (1988), however, studied 159 adult incest offenders and found their sexual offenses frequently extended beyond the family. Forty-nine percent had histories of being involved in nonincestual female pedophilia, 12% in male pedophilia, and 19% in rape.

THE CYCLE OF VIOLENCE

The cycle of violence is passed from one generation to the next. Early childhood victimization increases the risk for subsequent delinquent behavior, violent behavior and criminality (Widom, 1989, Kobayashi et al., 1995, Hunter and Becker, 1999). There is often a significant history of child maltreatment including neglect, physical and sexual abuse in the early lives of juvenile sex offenders (Knight and Prentky 1993; Ryan et al., 1996). Kobayashi et al. (1995) found that physical abuse by fathers and sexual abuse by males increased sexual aggression by adolescents. While the exposure of juvenile sex offenders to physical violence is comparable to that generally found in other delinquents emphasis has been placed on the increased reports of sexual victimization in the history of sexual abusers.

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

Sexual behavior is commonly affected by a history of sexual victimization. Boys and girls exposed to sexual abuse and deviant sexual experiences, behaviors, attitudes and knowledge are at risk for early erotization and precocious sexualization. There is some evidence that boys are more likely to be sexually aroused at the time of sexual victimization and to subsequently exhibit more sexual behaviors when compared to sexually abused girls (Friedrich, 1995; McClellan et al., 1997).

A history of sexual abuse is more prevalent in sexual abusers than in the general population and in nonsexual abusers (Dhawan and Marshall, 1996; Friedrich, 1995). Reports of sexual victimization in the history of adolescent sex offenders vary from 19 to 82% (Becker et al., 1986; Dhawan and Marshall, 1996; Fehrenbach et al., 1986; Longo, 1982; Kahn and Chambers, 1991; Ryan et al., 1996; Shaw et al., 1993, Zgourides et al., 1994). The victims of sexual abuse may internalize the aggressivized and erotized facets of the sexual experiences into preferred patterns of deviant sexual gratification through a process of social learning, imitation, modeling and identificatory pathways. Becker et al. (1989, 1992) demonstrated that a history of physical and sexual abuse in juvenile sex offenders is associated with higher phallometrically measured arousal to both deviant and non-deviant sexual stimuli. She found that adolescents who had been sexually victimized manifested more deviant erectile responses than adolescents who had not been sexually abused. Boys who have been sexually abused often demonstrate higher rates of externalized behaviors such as aggression, impulsivity and temper tantrums as well as sexual preoccupations and sexually inappropriate behaviors (Holmes and Slap, 1998). Cooper et al. (1996) compared sexually abused and non-abused adolescent sexual abusers and found that the abused sexual abuser had an earlier onset of sexual offending behavior, had more victims, and was more likely to have both male and female victims and to manifest greater psychopathology and interpersonal problems.

Early sexual sensitizing experiences may determine deviant patterns of erotization. Developmental experiences with significant others are crucial to the elaboration of a sexual life. It is known that animate or inanimate non-sexual arousing stimuli may through pairing with sexual-arousing stimuli acquire a conditioned sexually evocative status. Patterns of sexual arousal may become coincidentally paired with aggressive sexuality, enemas, corporal punishment, abusive discipline or other accidental and sometimes terrifying experiences. These early experiences if associated with sexual arousal and sexual excitement may become woven into the preferred patterns of sexual gratification. Boys and girls who have been sexually victimized may experience their first sexual arousal during the victimization. The factors related to and derived from the sexual abuse experience which are thought to be associated with increased risk for inappropriate sexual behaviors are sexual arousal at the time of the sexual

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

abuse; uncertainty and confusion about sexual identity; compensatory hypermasculinity; and a readiness to reenact the sexually victimizing experience (Friedrich, 1995; Watkins and Bentovim, 1992).

There is increasing awareness of the importance of early sexual abusive experience in precipitating inappropriate sexual behaviors in very young children (Holmes and Slap, 1998). McClellan et al. (1996) in a retrospective chart review of 499 mentally ill youth (5 to 18 years) found that the probability of engaging in any sexually inappropriate behavior was inversely related to the age they were sexually abused. A history of sexual abuse before 7 years of age was significantly associated with hypersexuality, exposing and victimizing sexual behaviors. A number of studies have found that sexually abused children may become sexually reactive and are more sexually inappropriate and manifest higher ratings on sexual activities scales (Deblinger et al., 1989; Friedrich, 1993). Yates (1982) described sexually victimized children, 2 to 6 years, who were found to exhibit excessive erotization. These children had trouble differentiating sensual from affectionate touching and erotic from non-erotic relationships. They were easily sexually aroused and quick to turn to others for sexual gratification. Kendall-Tackett et al. (1993, p. 165) noted that children who had been sexually abused were more likely to exhibit sexualized behaviors such as “sexualized play with dolls, putting objects into the anuses or vaginas, excessive or public masturbation, seductive behavior, requesting sexual stimulation and age inappropriate sexual knowledge.”

It is known that the younger the child when he commits his first sexual offense the more likely that child has been sexually victimized. (Johnson, 1988; McClellan et al., 1996). Johnson (1988) found that 72% of child sex offenders under six years of age had been sexually victimized; 42% of those 7-10 years of age; and 35% of those 11-12 years. Shaw et al. (1993, 1996) documented the high prevalence of early sexual victimization in the history of adolescent sex offenders (67%) who had been committed to a residential center. They found that the mean age of sexual victimization was 6 years of age with 35% of the group having been sexually abused before 5 years of age. While 30% of the adolescent sex offenders were not directly sexually abused, they invariably came from homes where they were prematurely exposed to sexual violence, promiscuity and pornography. Ford and Linney (1995) observed that juvenile sex offenders had been exposed to pornographic materials and/or “hard core” sex magazines at an earlier age than non-sexual violent offenders. Longo (1982) found that among his adolescent sex offenders there was a higher than expected history of direct consenting sexual

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

contact with older males and females. These adolescents reported feeling uncomfortable, insecure, inexperienced, inferior and unable to satisfy their older sex partners.

As important as sexual victimization is in the history of juvenile sexual abusers, it is not explanatory. Most victims of sexual abuse do not grow up to be abusers. As a form of child maltreatment it is only one of a number of critical factors which may contribute to the risk of becoming sexually abusive (Prentky et al., 1997). Nevertheless it is essential to understand the role of sexual victimization in the patterning of sexual offending behavior and as a critical factor in the planning of therapeutic interventions.

CLINICAL PRESENTATION

Early studies of inappropriate sexual behavior in juveniles were concerned with trying to define a child molester syndrome or profile (Schoor et al., 1966). The complex multidetermined nature of juvenile sexually aggressive behavior has made it difficult to set up a predictable taxonomic system. Presently, there is no evidence that any one profile or typology is characteristic of juvenile sex offenders (Becker and Hunter, 1993; Levin and Stava, 1987).

Adolescents who display sexually abusive behavior are a heterogeneous population. They are represented in every social economic class, and racial, ethnic, religious, and cultural group. There are, however, a number of factors that have been commonly found in the history of adolescent sex offenders. These include: impaired social and interpersonal skills; prior delinquent behavior; impulsivity; academic and school problems; family instability; family violence; abuse and neglect; and psychopathology (Araji, 1997; Becker and Hunter, 1993; Fehrenbach et al., 1986; Ryan et al., 1996).

THE PREADOLESCENT SEX ABUSER

There is increasing awareness that children under 12 years of age may be sexually abusive toward other children (Araji, 1997; Gray et. al., 1997; Johnson, 1988, 1989, 1999). In these cases it is necessary to discriminate between age appropriate sexual exploration and sexually abusive behavior. Appropriate sexual exploratory behavior is usually carried out with children of the same age and size, and with mutual consent.

The age differential between the abuser and the victim in children is less definitive. It may be less than two years and in some instances the younger child may be the initiator of sexual

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

behaviors. For those children 9 years and older the age difference between abuser and victim is usually greater than two years.

Younger sexual abusers are often reacting to their own histories of sexual victimization (Gil and Johnson, 1993; Johnson 1988). The term “sexually reactive” refers to children who display sexually inappropriate behavior in response to sexual abuse or exposure to explicit sexual stimuli (Gil and Johnson, 1993). They have been characterized as often exhibiting impulsivity, anger, fear, loneliness, confusion, depression, obsessional and compulsive preoccupation with sex, excessive sexualization, anxiety and sleep disturbance (Araji, 1997; Johnson and Berry, 1989). In approximately half of the cases, the sexual victims of these children are their own siblings. The sexual abusive behaviors between children are usually characterized by themes of secrecy, dominance, coercion, threat and force. The sexual behaviors are far advanced beyond those expected for the age of the child and may include oral and vaginal intercourse and forcible penetration of the anus or vagina with fingers or other objects. The sexual behaviors are patterned, increase over time, and are often associated with conduct disordered behaviors and may continue in spite of intervention (Araji, 1997; Gil and Johnson, 1993).

A study of 6 to 12 year old children who had exhibited sexual misbehaviors found a number of characteristics indicative of parental and family distress, i.e., violence between parents, history of sexual victimization and perpetration within the extended family, physical abuse, poverty, the witnessing of domestic violence, parental arrests, and the prior use of educational and therapeutic services for children (Gray et. al., 1997). Younger children exhibiting sexual misbehaviors are more likely to have been sexually and physically abused at a younger age and to manifest more problematic sexual behaviors with a higher percentage of hands on sexual behaviors.

THE ADOLESCENT SEX ABUSER

Becker (1988a) has suggested that there are essentially 4 kinds of sexual abusers with most offenders combining features of each:

- The true paraphiliac with a well-established deviant pattern of sexual arousal
- The antisocial youth whose sexual offending behavior is but one facet of his opportunistically exploiting others

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

- The adolescent compromised by a psychiatric or neurologic biological substrate disorder which interferes with his ability to regulate and modulate aggressive and sexual impulses
- The youth whose impaired social and interpersonal skills results in his turning to younger children for sexual gratification unavailable from his peer groups.

Sexually aggressive adolescents are more likely to endorse traditional sex-role stereotypes, male dominance, and rape-supportive myths and to have negative and stereotypical attitudes about women (Epps et al., 1994; Segal and Stermac, 1984; White and Koss, 1993). The history of prior sexual offenses at the time of apprehension varies from 8 to 65% depending on if the history was obtained from self-report, prior offense records, and the specific sample of sexual abusers studied (Fehrenbach et al., 1986; Ryan et al., 1996; Shaw et al., 1996; Smith and Monastersky, 1986).

PSYCHOPATHOLOGY

Sexual offending behavior is associated with a matrix of behavioral, emotional and developmental problems (Awad and Saunders, 1989; Kavoussi et al., 1988; Shaw et al., 1993; 1996). The studies of the psychological characteristics of sexual abusers, however, may be skewed as only a small percentage of sexual abusers are actually caught (Harvey and Herman, 1992).

Investigators have examined adolescent sexual abusers in an effort to explicate the spectrum of psychopathology (Becker, 1988a; Lewis et al., 1979; Kavoussi et al., 1988; Shaw et al., 1993; Shaw et al., 1996). The findings are understandably affected by the assessment procedures, diagnostic instruments, and sample selection, i.e., inpatient-outpatient and severity of offending behavior.

When adolescent sexual abusers are compared to psychiatric controls or normals on the MMPI they manifest more psychopathology, demonstrate more global maladjustment, and are more threatened by heterosexual interactions (Herkov et al., 1996; Katz, 1990).

Studies comparing violent and non-violent delinquents with juvenile sexual abusers across a broad range of intellectual, psychological, neuropsychological, and psychoeducational variables have found few significant differences between the groups (Lewis et al., 1979; Tartar et al., 1983; Hastings et al., 1997; Jacobs et al., 1997). Truscott (1993) compared sexual, violent and property offenders and noted that the sex offenders were twice as likely to have a history of being sexually abused. Juvenile sex offenders generally do less well on academic achievement

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

tests than nonsexual delinquents and conduct disordered youth (Lewis et al., 1979; Shaw, et al., 1993).

A number of studies have focused on the specific mental disorders found among juvenile sex offenders. Psychiatric co-morbidity has been found in approximately 60-90% of adolescent sexual abusers. The most prevalent co-morbid psychiatric disorders are conduct disorder, 45-80%; mood disorders, 35-50%; anxiety disorders, 30-50%; substance abuse, 20-30%; and attention-deficit hyperactivity disorder, 10-20% (Becker et al., 1986, 1991; Kavoussi et al., 1988; Shaw et al., 1993). The younger the child when he committed his first sexual offense the higher the number of coexisting psychiatric diagnoses (Shaw et al., 1996).

DELINQUENCY

Reports of prior nonsexual delinquent behavior in the history of juvenile sexual abusers varies from 27-91% (Awad et al., 1989; Becker et al., 1986; Fehrenbach et al., 1986; Ryan et al. 1996; Saunders et al., 1986; Schoor et al., 1966; Shaw et al., 1993). A national survey of juvenile sex offenders reported that 63% had previously committed nonsexual delinquent offenses (Ryan et al., 1996). Twenty-eight percent had a history of committing 3 or more nonsexual criminal offenses.

Elliott (1994b) in the National Youth Survey, a longitudinal study of a national probability sample of 1,725 youths, 11-17 years, reported that rape is the final step in an escalating sequence of violent criminal activity. The usual sequence is of criminal behavior evolving from aggravated assault to robbery to rape. The peak age for males to commit a sexual assault was 17 years of age. Rapists were noted to have committed virtually every form of violent offenses. It is evident that there is a subset of sexual-aggressive offenders characterized by core antisocial features in which the sexual aggression is only one facet of a life style in which the individual opportunistically exploits others for personal gain (Ageton, 1983; Lewis et al., 1979, Shaw et al., 1993; Hastings et. al., 1997).

PERSONALITY CHARACTERISTICS

Adult sex offenders frequently manifest significant personality disturbances (Packard and Rosner, 1985). Shaw et al. (1996) in an effort to explicate personality traits and character pathology examined a population of adolescent sex offenders in a residential center using a structured psychiatric interview. He found a high prevalence of severe personality traits to

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

include narcissistic, borderline and conduct disordered behaviors. Ninety-two percent of the boys met criteria for a diagnosis of conduct disorder; 67%, narcissistic personality disorder; and 72%, borderline personality disorder. The younger the age of onset of sexually offending behavior and the younger the sex offender was at the time of his own sexual victimization the more likely he was to exhibit symptoms of a borderline personality. The high prevalence of narcissistic and borderline psychopathology is consistent with histories of severe emotional, physical and sexual abuse. Carpenter et al. (1995) found that adolescent sex abusers who offended against children were more schizoid, dependent, and avoidant than those adolescents who sexually abused peers.

FAMILY ENVIRONMENT

Most juvenile sex offenders live at home at the time of committing their sexual offenses (Kahn and Chambers, 1991; Ryan et al., 1996). The family environment is usually characterized by family conflict, family instability, family dysfunction, and exposure to violence (Awad et al., 1984; Becker et al., 1993); harsh inconsistent parenting, and physical and sexual maltreatment (Becker et al., 1993; Knight and Prentky, 1993; Lewis et al., 1979; Shaw et al., 1993); and low adaptability and low cohesion (Smith and Monastersky, 1986).

SOCIAL AND INTERPERSONAL SKILLS

Adolescent sex offenders manifest impaired social and interpersonal skills. Fehrenbach and Monastersky (1988) found that 65% were socially isolated and that 32% did not have a friend. Awad et al. (1984) reported that 46% of his sample of adolescent sex offenders were loners. There is some evidence that adolescent sex offenders unable to relate to their own peer group frequently turn to younger children for the gratifications denied to them by their own peer group (Awad and Saunders, 1989; Schoor et al., 1966).

SCHOOL AND ACADEMIC PROBLEMS

Juvenile sex offenders usually present with a history of academic and school behavior problems and perform less well on tests of academic skills (Awad et al., 1984; Awad and Saunders 1989; Fehrenbach and Monastersky, 1988; Lewis et al., 1979; Ryan et al., 1996; Shaw et al., 1993). Studies have found that approximately 40-80% of juvenile sex offenders manifest

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

learning disabilities and behavior problems in school (Awad and Saunders, 1989; Ryan et al., 1996; Shaw et al., 1993).

THE FEMALE JUVENILE SEX ABUSER

There have been few studies of female sexual aggression (Fehrenbach and Monastersky, 1988; Green, 1999; Green and Kaplan, 1994; Knopp and Lackey, 1987; Kaplan and Green 1995; Mathews et al., 1997). Anderson (1996) surveyed young college women and found that 29% had engaged in sexual coercion; 28% had committed sexual abuse; and that 43% reported initiating sexual contact by using sexual aggressive strategies. It is known, however, that 50 to 95% of female sexual abusers report a history of sexual victimization (Kaplan and Green, 1995; Green, 1999). Female juvenile offenders are more likely to have been sexually abused at a younger age and to have had multiple abusers and were three times more likely to have been sexually abused by a female (Mathews et al., 1997).

Fehrenbach and Monastersky (1988) studied 28 female adolescent sex offenders, 10-18 years of age, with a mean age of 13.6 years. Fifteen had been charged with rape and 13 with taking of indecent liberties. Ten of the female sexual abusers had assaulted males, 16 had assaulted females and 2 had assaulted both males and females. Approximately 70% of the offenses took place while baby sitting. The mean age of the victims was 5.2 years. Fourteen (50%) of the offenders had a history of being sexually abused.

Mathews et al. (1989) have described 5 variations commonly found among adult female sex abusers: 1) female adolescents who opportunistically exploit and fondle younger children; 2) females with a history of being sexually abused who subsequently abuse younger children; 3) an older female who falls in love with a younger child that she is mentoring; 4) emotionally unstable females with an impaired capacity to modulate and regulate their own sexual impulses; and, 5) females who sexually abuse children only in the company of a male sexual abuser.

THE DEVELOPMENTALLY DISABLED JUVENILE SEX ABUSER

There is some suggestion that developmental disabilities are over-represented in juvenile sex offenders. The prevalence of sexual offending behavior is at least as common if not more common in the disabled population (Gilby et al., 1989; Stermac and Sheridan, 1993). McCurry et al. (1998) in a retrospective chart review of 200 seriously mentally ill youth (5 to 18 years) found that 50% had engaged in problem sexual behaviors and that independent of a history of sexual abuse low verbal IQ was correlated with sexually inappropriate behavior. Twenty (65%)

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

of those with FSIQ (WISC-R/WAIS-R) IQs less than 70 engaged in inappropriate sexual behaviors.

The clinical characteristics of the disabled sex offenders, the spectrum of sexual offenses and the victim profiles are not demonstratively different from the nondisabled sex offender (Stermac and Sheridan, 1993). The paucity of research studies limits our overall understanding of this group of sexual abuser. There is recognition that treatment strategies have to be tailored to their cognitive limitations and are usually more focused on behavioral interventions (Murphy et al., 1983).

ASSESSMENT

The assessment of sexually abusive youth is carried out for the purpose of understanding the sexual abuser so that one may make reasonable judgments regarding the protection of the community, case disposition, and treatment planning (Dougher, 1988; Ross and Loss, 1991). The clinical assessment of juvenile sex abusers requires the same comprehensive evaluation as other children and adolescents (American Academy of Child and Adolescent Psychiatry, 1997). It necessarily requires the use of multiple informants and multiple sources of data. Important sources of information are medical and psychological reports, offense reports, victim statements, protective services reports, and probation reports. Ideally the collateral information should be obtained before the individual interview; otherwise, one is left relatively helpless before the offender's normal proclivity to minimize and deny. There are no empirical measures or psychological batteries that can substitute for a careful assessment.

FORENSIC CONSIDERATIONS

It is essential that the clinician define his role in the assessment of the sexual abuser. While many of the issues are relevant to a forensic evaluation, these parameters are designed to provide guidelines for the clinical evaluation of the sexual abuser. It is generally preferable to conduct the clinical evaluation after adjudication. The focus of the clinical interview is on assessing amenability to treatment, required levels of care, treatment goals and objectives and risk of reoffending rather than a determination of guilt or innocence (Hunter and Lexier, 1999).

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

The juvenile sex abuser is advised of reporting laws and the limits of professional confidentiality. An informed consent signed by the juvenile and his parent/guardian should be obtained prior to the clinical interview. It is important to educate and clarify for the individual and his family what is going to happen and when. The role of protective services and the juvenile justice system should be explained when relevant. Consent forms should be developed to cover the use of controversial assessment and treatment procedures such as phallometric assessment, aversive conditioning and medications that are not accepted as standard of practice. The role of the protective services and the juvenile justice system may have to be explained. It is important to educate and clarify for the individual and his family what is going to happen and when.

One of the difficulties in evaluating sexually offending youth is the contradictory intentions of those who provide clinical services and those representatives of the court who are interested in prosecution or defense. Information obtained by the clinician during the clinical interview may be subpoenaed by the court. The clinician may have to testify under oath as to what he knows about the sexual history and activities of the offender. In some instances youth will be told by their attorney not to report any of their sexual history or activities to the clinician because of impending legal and court procedures. In these instances only a partial and limited sexual history will be obtained, and it will be necessary to wait until the court has determined responsibility before one can complete the evaluation.

THE CLINICAL INTERVIEW

The cornerstone of the assessment and evaluation of the sexual abusers is an extensive and comprehensive individual clinical interview. It is common for sexual abusers to deny, minimize, and disavow their sexual offenses. Sefarbi (1990) found that half of adolescent sexual abusers initially denied the sexual offending behavior at the time of clinical referral and that this denial was usually supported by the family. Ryan et al. (1996) in her national survey of juvenile sex offenders found that at the time of the evaluation only 19.4% accepted full responsibility for the sexual offense. Thirty-three percent admitted little or no responsibility. Sixty-two percent expressed little or no empathy for the victim. Fifty-one percent expressed little or no remorse. Although nearly two-thirds accepted blame for the sexual offense, one-third blamed the victims.

The denials usually take the form of protestations of innocence, statements that they are wrongly accused, or overt attempts to discount the victim's credibility. Not uncommonly, the offender states the victim consented to the sexual activity in spite of all the evidence to the

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

contrary. Considerable time and effort will be necessary to unravel the underlying motivations and sexual fantasies.

It is necessary to establish the nature of the sexually abusive behavior. This is often a difficult task because of the interplay of a number of variables which may impede the recovery of the exact details of what transpired. Since frequently laws have been transgressed the offender is often less than forthcoming. Issues of shame, guilt and fear of punishment impede disclosure. Most offenders are not motivated to disclose the circumstances of their sexually abusive behavior until they are confronted or fear the consequences if they do not talk. It is essential that the clinician adopt a nonjudgmental stance and relates to the juvenile offender in a matter of fact exploratory manner. While the clinician has a structured agenda it is advisable to follow the patient's lead and take advantage of open-ended questions if there are promising opportunities. The clinician avoids sexual jargon, maintains a sense of seriousness, and avoids any minimization of the seriousness of the sexually abusive behavior. Generally it is best to initiate the interview with a nonthreatening line of questioning and thereby avoid the initial defensiveness. The interviewer develops lines of questioning that will allow him/her to know more about the offender, his family, school and his current life situation. Confront minimization, denial, and the omissions of important information. There is little value in getting angry and accusatory. It is more useful to be patient, persistent, and not easily dissuaded.

The Assessment of the Sexual Abusive Behavior

In the evaluation the clinician is interested in answering a number of questions regarding the abuser and the sexual abusive incident (see Table 2) (Groth et al., 1981; Ross and Loss, 1991; Saunders and Awad, 1988).

Insert Table 2 Here

Sexual History

As part of the ongoing psychiatric assessment it is essential to obtain a comprehensive sexual history. This is best accomplished after there has been some time to develop a trusting relationship. It works best to have a conceptual framework and sequence of questions which directs the inquiry.

In the evaluation, the interviewer determines the child's sexual knowledge and education, sexual development, and sexual experiences. Inquiries are directed to learn what the child

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

knows about gender differences, sexual intercourse, auto-erotic practices, and his/her preferred patterns of sexual behaviors. Specific questions may be asked regarding the child's understanding and knowledge of normal sexual activities such as kissing, dating, petting, masturbation, whether he/she has been sexually active and engaged in intercourse, homoerotic experiences, etc. It is necessary to discriminate between compulsive sexual behaviors and paraphiliac compulsive sexual behaviors. In the former condition the individual displays conventional and normative sexual behaviors but they are excessive and carried to the extreme (Coleman, 1990, 1992). It has been suggested that non-paraphiliac compulsive sexual behaviors may be a variant of obsessive-compulsive disorder.

The clinician will need to explore the history of sexual aggressive behaviors including: previous incidents of sexual abusive behavior, the pattern and spectrum of previously committed sexual aggressive acts; the victim profile; the internal and external triggers that initiate the sexual abuse cycle; the role of aggression and sadism in the sexual offense; the need to dominate, control and humiliate the victim; the erotization of the aggression; the history of sexual victimization, physical abuse and emotional neglect; the history of exposure to inappropriate and sexually explicit behavior; and the history of prior non-sexual delinquent behavior.

Developmental and Psychosocial History

Other areas of the assessment process are those associated with a comprehensive developmental history, i.e., the nature of the pregnancy, perinatal history, developmental milestones, family relationships, early identificatory models, capacity for relationships, school experiences, social skills, substance abuse, and prior medical and psychiatric history. The family assessment provides an opportunity to understand the early developmental and environmental context within which the sexual abuser developed. Information is obtained regarding the parents' personal and psychological history, their use of authority and discipline, and the role of coercive sexuality in the family. How is affection, tenderness, competition, aggression, love, sexuality and lust expressed in the family? How supportive and available is the family as a treatment resource?

Legal History

Is there a history of arrests, convictions, incarcerations, use of weapons, or cruelty to animals?

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

Medical and Psychiatric History

It is important to obtain a comprehensive medical and psychiatric history with specific attention to sexually transmitted diseases, HIV infection, psychopathology and psychiatric comorbidity.

School and Academic History

A specific area of concern is the evaluation of intellectual capacities and academic performance. Information is obtained from the school and from formal psychoeducational and psychological assessments.

Mental Status Examination

A comprehensive mental status examination is carried out to assess the presence of psychopathology, personality disturbances, organicity, substance abuse and to acquire an understanding of adaptive, coping and defensive strategies. Suicidal content and risk should be assessed specifically. Apprehension by judicial authority and the associated shame of exposure, embarrassment, stigmatization, fear of punishment and incarceration are risk factors for suicidal behavior. A careful assessment of the spectrum of suicidal behavior is undertaken to establish the degree to which suicidal ideation, history of suicidal behaviors, threats or plans are present.

PSYCHOLOGICAL TESTING

There are no specific empirical measures or psychometric tests which can identify, diagnose, or classify sexual abusers. Psychological tests are used adjunctively as part of an overall comprehensive evaluation to understand the personality, motivations, ego strengths, intelligence, defense and coping strategies, psychopathology, sexual knowledge and sexual behaviors of the offender. The most commonly used psychological measures are the Minnesota Multiphasic Personality Inventory (MMPI); WISC-III or WAIS; Rorschach; the Thematic Apperception Test; the Bender Gestalt; and Draw-a-Person. While projective tests have been historically important, their use is limited to what they may contribute to the comprehensive assessment procedure. Neuropsychological testing and psychoeducational assessment may be required when one is suspicious of neurologically based deficits and/or learning disabilities. Measures of learning are an essential part of the assessment procedure. When indicated, family assessment measures may be administered to more fully unravel family dynamics and family process.

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

Specific tests which have been used with adolescent sex offenders are the following:

The Multiphasic Sex Inventory

The Multiphasic Sex Inventory (MSI) developed by Nichols and Molinder (1984) has been standardized for adolescents. The latter measure is a 300-item, true-false test with 14 clinical and validity scales which provide measures regarding sexually offensive behavior, sexual deviations, sexual dysfunction, sexual knowledge and sexual attitudes. There is a lack of published data on the reliability and the validity of this instrument in the adolescent population (Becker and Hunter, 1997).

The Adolescent Cognitions Scale

The Adolescent Cognitions Scale for Juvenile Sex Offenders, a 32-item, forced-choice inventory, was found by Hunter et al. (1991) to have only marginal reliability and failed to discriminate between sexual and nonsexual abusers.

The Adolescent Sexual Interest Card Sort

The Adolescent Sexual Interest Card Sort, a 64-item, self report measure of sexual interest is reported to correlate poorly with phallometric assessments of sexual interest. Hunter et al. (1995) suggest caution in its use.

Child Sexual Behavior Inventory

The Child Sexual Behavior Inventory (CSBI) is a measure that utilizes parent report of sexual behavior in children ages 2-12 years of age and provides information on a number of domains to include sexual anxiety, sexual interests, sexual knowledge, sexual intrusiveness, gender role behavior, boundary problems, etc. (Friedrich, 1997). While increasingly used, it is limited by its reliance on parent report.

PHALLOMETRIC ASSESSMENT

Some authors have suggested the use of phallometric testing, the measuring of penile erection in response to different stimuli, as a way to determine sexual preferences (Freund, 1963). Prentky et al.(1997) noted that phallometric assessment of sexual arousal in response to depictions of children can differentiate child molesters from nonmolesters, same-sex molesters

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

from opposite sex molesters, and extrafamilial molesters from incest offenders. This technique is usually reserved for the most severe and repeat sexual abusers. This procedure has generally been used with caution with minors cause of the lack of empirical studies, problems of obtaining informed consent, and a reluctance to expose children and adolescents to further sexual stimulation through the portrayal of deviant sexual activities (Saunders and Awad, 1988).

Becker and Hunter (1997) report that the relationship between phallometric arousal and certain clinical characteristics in the adolescent population are weaker than in the adult population. One study of adolescents has suggested caution in the use of phallometric measures as there is an age effect determining sexual arousal and erectile measures (Kaemingk et. al., 1995).

DISPOSITION

At the end of the assessment process, the clinician should be prepared to address the following issues and provide guidance to other professionals, the juvenile court and other community agencies.

- The degree of the abuser's threat to the community, victim or other potential victims
- The risk of repeating the sexually aggressive behavior
- The treatment needs of the individual and his family
- The desirability of removing the abuser from his/her family
- The appropriate treatment program for the abuser (i.e. a community outpatient program) or whether he/she needs to be placed in a more restrictive environment (i.e., detention center, residential program)

TREATMENT PLANNING

There is increasing awareness that sexually abusive behavior is more than a simple disorder of sexual arousal (Schwartz, 1992). There is a need for combined and integrated treatment approaches (Becker, 1994). The spectrum of emotional, behavioral, and developmental problems presented by these young people requires an integrated multi-model treatment program which is tailored to the individual's clinical presentation, social and family support system. The predominant treatment approaches include cognitive-behavioral and psycho-educational modules, behavioral interventions, relapse prevention, psychosocial

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

therapies, and psychopharmacological approaches (Borduin et al., 1990; Becker, 1988a, 1994; Becker and Hunter, 1993; Schwartz, 1992; Shaw, 1999).

Treatment planning necessarily reflects a judgment about the level of care required. Decision making is complicated by the heterogeneity of this population. Sexual abusers vary as to the severity of delinquency/criminality, psychopathology, sexual deviancy, the history of child maltreatment and sexual victimization, family support systems, motivation for help and the danger they represent to the community. Hunter and Figueredo (1999) found that the degree of sexual maladjustment, denial, and the lack of a sense of accountability for one's sexual offenses predicted failure in treatment compliance in a community based intervention program. Sexually reactive children reenacting their own sexual victimization and erotization often respond well to individual and family interventions.

More serious sexual abusive behavior is usually embedded in an array of developmental, behavioral and emotional problems. Most juvenile sexual abusers, however, do not meet diagnostic criteria for paraphilia nor do they have well established patterns of deviant sexual arousal. The youthful nature of the offenders suggests that in many instances the offenders are still experimenting with different sexual activities. The plasticity of their cognitively maturing structures and the ongoing process of internalizing social and interpersonal skills suggest considerable malleability and potential for responding to appropriate social learning and treatment.

Juvenile sexual abusers usually present with considerable psychiatric co-morbidity. A number of studies have documented the frequency and diversity of the psychiatric impairment (Becker et al., 1991; Kavoussi et al., 1988; Lewis et al., 1979; Shaw et al., 1993, 1996, 1999). Psychiatric diagnoses of conduct disorder, depression, anxiety, and substance abuse are common as well as evidence of character pathology and personality disorders. An understanding of the psychopathology and personality structure of juvenile sex offenders is necessary to plan a comprehensive treatment program.

An essential element in treatment planning is the evaluation of the severity of the sexual offending behavior and the risk of recurrence of sexual offending behavior. This is a difficult task even when the judgements are made by experienced clinicians (Kahn and Chambers, 1991). Since the majority of adolescent sex offenders have a prior history of nonsexual delinquent acts, the clinician is concerned about recidivism not only for the sexual offenses but for nonsexual delinquent acts. There is evidence that the recidivism is higher for nonsexual delinquent acts than it is for the sexual offenses.

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

The determination of the risk of reoffending is derived from the extensive clinical, developmental and historical assessment of the offending youth (Gerdes et al., 1995, Rasmussen, 1999). Considerations in evaluating the risk of further sexual offenses include: the frequency and diversity of the sexual offenses; the severity of the aggressive-sadistic behavior; the planfulness/impulsivity of the sexual offending behavior; psychopathology; neurological impairment; prior antisocial or violent behavior; motivation for treatment; intelligence; psychological mindedness; capacity for empathy and, family, community, and social support.

Factors to consider for placement in a more restrictive environment include: the consistent need to deny sexual offenses; the lack of remorse and victim empathy; a well established pattern of frequent and diverse committed acts of sexual aggression; the number of previous arrests; the number of victims; severity of psychopathology; failure of previous treatment efforts; the degree of compulsivity and sexual arousal; and a documented history of violent aggressive and sadistic behavior (Hunter and Figueredo, 1999; Rasmussen, 1999).

In the final stages of the clinical interview and assessment procedure it is helpful to educate and clarify for the individual and his/her family what is going to happen and when. The role of protective services and the juvenile justice system may have to be explained. Provide realistic assurance that his/her problems are treatable, that he/she can learn to control and regulate sexual and aggressive impulses and that there are specific treatment programs for children and adolescents with these problems. Explain to the family members what their participation will be in the treatment program. It is necessary to explain your recommendations to the offender and what is generally expected of the offender and his family in treatment.

Traditionally, treatability has been related to the sexual abuser's willingness to accept accountability for his sexual offenses manifested by: 1) admission of his sexual offense; 2) acceptance of his sexual offending behavior as a problem; 3) motivation stop sexual offending; and, 4) willingness to participate fully in treatment. Other reports suggest some success with adult sexual abusers who initially denied their sexually abusive behavior (Schlank and Shaw 1996; Maletzky 1996). The variables that seemed to be associated with sexual recidivism are a readiness to blame the victims, the number of victims, history of sexual abuse, and the use of threats in the commission of the sexual crime (Kahn and Chambers, 1991; Rasmussen, 1999).

TREATMENT

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

The first task is to protect the community. It is a social imperative that the population of adolescent sex offenders who are at risk to be the adult sex offenders of tomorrow be the focus of prevention-intervention strategies. Prentky and Burgess (1990) in a cost-effective analysis of sex offender treatment programs noted a substantial savings to the community not only in terms of cost but also in sexual reoffending behavior. Farrell and O'Brien (1988) found that the cost of providing specialized outpatient treatment was about one-fifteenth of the cost that one would have to provide in an institutional or residential setting. A significant percentage of juvenile sexual abusers will respond to therapeutic intervention (Becker, 1994; Becker and Hunter, 1997; Bremer, 1992; Dwyer, 1997; Hall, 1995).

Little is known about the natural course of untreated juvenile sex offending behavior. Dorshay (1943) found very little recidivism in a 6-year follow up study of 256 juvenile sex offenders. He found that 7% reoffended sexually while 40% committed other nonsexual delinquent acts. Marshall and Barbaree (1988) have estimated that the recidivism rate for untreated sexual abusers is 40%. Rubenstein et al. (1993) found in a small sample of severe adolescent sex offenders that 37% had sexually re-offended and that 89% continued to commit other acts of violence.

There is general agreement that the recidivism rate for treatment programs is in the range of 5-15% (Becker, 1990; Becker and Hunter, 1993; Kahn and Chambers, 1991; Knopp, 1991; Knopp et al., 1992; Smith and Monastersky, 1986). Questions remain, in spite of the number of studies, suggesting a low recidivism rate among treated juvenile sex offenders. Critical commentary of treatment studies have focused on methodological flaws; specifically the lack of random assignment, controls, and post test design only paradigms (Furby et al. 1989; Miner 1997). Maletzky (1997) has suggested, however, that few would argue for untreated control groups of juvenile sex offenders. It is unethical to withhold treatment for those who want treatment and unethical to fail to protect the community by failing to intervene with juvenile sex offenders. In spite of the lack of empirical rigor, there is considerable evidence that treatment interventions are effective in interrupting the course of sexual abusive behavior. Hall (1995) in a metaanalysis of 12 studies found a small but significant effect of treatment over no treatment.

There are several explanations as to why the adolescent offender may be more amenable to treatment than the adult sex offender:

- The adolescent offender's deviant pattern of sexual offending behavior is less deeply ingrained
- The adolescent is still exploring alternative pathways to sexual gratification

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

- The adolescent's central masturbatory fantasy is still evolving and is not fully consolidated
- The adolescent is available for learning more effective interpersonal and social skills

Despite the successes indicated in these studies it is evident that not all adolescent sex offenders are equally treatable. Treatability depends on a number of patient characteristics and situational factors: the level of understanding of the seriousness of the offense; the motivation to discuss and understand the offense; the capacity for empathy and human relatedness; the severity of psychopathology; the entrenchment of deviant sexual arousal patterns; the type and frequency of sexual offending behavior; the aggressivization of the sexual offense; the degree of characterological impairment; and the nature of the treatment program (Groth et al., 1981, Shaw et al., 1996; Smith and Monastersky, 1986).

The treatment of juvenile sex offender has generally focused on a number of goals (Becker and Hunter, 1997; Becker, 1994; Ryan et al, 1987):

- Confronting the offender's denial
- Decreasing deviant sexual arousal
- Facilitating the development of non-deviant sexual interests
- Promoting victim empathy
- Enhancing social and interpersonal skills
- Assisting with values clarification
- Clarifying cognitive distortions
- Teaching the juvenile to recognize the internal and external antecedents of his sexual offending behavior

While treatment has been quite successful in reducing recidivism, adolescent sex offenders are not "cured." Treatment endeavors are organized to facilitate the sexual abuser's development of coping and adaptive strategies to prevent further sexual offenses.

The group setting is the preferred format for the treatment of sex offenders and is usually the medium through which cognitive-behavioral modalities, i.e., psycho-educational, behavioral and relapse prevention programs are conducted (Knopp et al., 1992; Schwartz, 1992; Smets and Cebula, 1987).

COGNITIVE-BEHAVIORAL INTERVENTIONS

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

Psycho-educational modules are didactic experiences that provide sexual abusers with information about sexuality, sexual deviancy, cognitive distortions, interpersonal and social behaviors, strategies for coping with aggressive and sexual impulses and anger management, etc. (Green, 1988, Becker and Hunter, 1997). This approach assumes that the offender has acquired a set of beliefs, attitudes, and expectancies which have shaped his/her sexual offending behavior and that his/her sexual behavior is maladaptive, contains “thinking errors,” and is associated with impaired communication and social skills (Johnston and Ward, 1996). These modules are taught by a therapist who often uses workbooks and homework assignments. The setting is usually a classroom although it may take place in the context of an ongoing group therapy. There is an emphasis on understanding the general patterns and determinants of sexual offending behavior, sex offender characteristics, and the spectrum of sexual offenses.

Specific Psycho-educational Modules

Victim Awareness/Empathy. The focus is on understanding the effects of sexual assault on the victim, identifying cognitive distortions and myths which support the sexual assault, and promoting participation in therapeutic endeavors.

Values Clarification. The therapist clarifies sexual values as they relate to the cessation of exploitative sexual relationships.

Cognitive Restructuring. This is an attempt to correct the cognitive distortions and the irrational beliefs which support the sexual offending behavior and to replace them with reality focused and culturally acceptable beliefs.

Anger Management. Instruction is provided to facilitate the recognition and the development of appropriate coping strategies for managing anger.

Assertiveness Training. Training is provided to promote more appropriate self assertive behavior to get one’s needs satisfied in reality oriented and culturally acceptable manner.

Social Skills Training. The therapist facilitates more effective prosocial behaviors, communication skills, and interpersonal awareness.

Sexual Education. The therapist provides information regarding human sexuality, myths, sex roles, and variations of sexual behaviors.

Stress Reduction/Relaxation Management. Techniques for coping and reducing stress, anxiety, and frustration are made available to the group

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

Autobiographical Awareness. Emphasis is on the individual developing an understanding of his/her own life trajectory and how the pattern of sexual offending behavior evolved over time.

Behavioral Interventions

Behavioral interventions have been used to diminish deviant sexual arousal and have been reported to be varyingly successful (Dougher, 1988). Some of the techniques are as follows:

Covert Sensitization. A counterconditioning paradigm in which the offender learns to extinguish pleasurable responses to sexually stimulating deviant imagery through the imagining of some negative reaction or aversive stimulus. Scenes are constructed for each offender according to his/her preferred sexual-erotic fantasies. (Cautela, 1966)

Assisted Covert Sensitization. Aversive stimuli such as noxious odors are used to facilitate an aversive reaction (Maletzky, 1974).

Imaginal Desensitization. The sex offender uses relaxation techniques to interrupt the sexually stimulating imagery and to inhibit the sexual arousal cycle (McConaghy et al., 1989).

Olfactory Conditioning. Sexually stimulating deviant imagery is presented which is followed by the presentation of a noxious odor.

Satiation Techniques. This involves either verbal or masturbatory satiation. The offender is first encouraged to masturbate to ejaculation in response to socially appropriate sexual fantasies with the concomitant feelings of affection and tenderness. Following this experience the offender is required to masturbate to deviant sexual fantasies. If he/she become aroused they are told to switch to an appropriate fantasy or in some instances exposed to an aversive stimulus such as ammonia (Gray, 1995). Verbal satiation requires the dictation on an audiotape of the most stimulating paraphiliac imagery for at least 30 minutes after masturbation 3 times a week. It is assumed that the paraphilic fantasy becomes boring and subsequently extinguished (Schwartz, 1992).

Sexual Arousal Reconditioning. This involves the pairing of sexual arousal with appropriate nondeviant sexual stimulation or sexual fantasies.

Relapse Prevention

Relapse prevention was originally developed as an intervention for substance abusers but was subsequently modified for sexual abusers (Pithers et al., 1983; 1988a; 1988b; Pithers and

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

Gray, 1996). Ninety percent of all sex offender treatment programs in North America report using relapse prevention (Pithers and Gray, 1996). This intervention strategy assumes that sexual offenses are not capricious happenings but are the product of contextual triggers and an array of emotional and cognitive precursors. The treatment process entails the explication and definition of each phase of the sexual assault cycle, i.e., the unique characteristics of each offender's cycle so that the offender will be aware of the triggers which initiate the cycle so that he will be alerted and employ new strategies for interrupting the sexual assault cycle (Ryan et al., 1987). Some of the emotional states which have been found to be important emotional triggers are boredom, social or sexual embarrassment, anger, fear of rejection, and numbness (Gray and Pithers, 1993). Proulx et al. (1996) found that "negative moods and conflicts" such as anger, loneliness, and humiliation coincided with deviant sexual fantasies and increased masturbatory behavior. The goals of relapse prevention are to empower the offender to manage his own sexual life through a cognitive understanding of the antecedents of his sexual offending behavior and through the development of coping strategies with which to interrupt the sexual offending cycle.

PSYCHOSOCIAL INTERVENTIONS

Interpersonal therapies include traditional individual approaches, family therapy, group therapies and the use of the therapeutic community.

Group Therapy

Group therapy with juvenile sex offenders provides a context in which the sexual abuser is unable to easily minimize, deny, or rationalize his sexual behaviors. Peer-related group therapy as the conduit for therapeutic interventions is used in a number of different ways depending on the setting (outpatient versus a more restrictive environment), group membership, the severity of the sexual offenses, group goals and objectives, whether the groups are open or closed, and the length of the group experience. Therapeutic community groups are often used in hospital or residential treatment settings as a vehicle for milieu administrative decision making and for the monitoring of a behavioral management system. Group membership is usually either all male or female and very rarely is there advantage in mixing sexes in a group designed for the treatment of sex offenders.

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

In group therapy, the sex offender is confronted by peers who are “street smart” and who are not easily manipulated and who are able to confront the attempts at minimization and denial. Denial may be very persistent as it is strongly supported by cognitive distortions and stereotypical views of females. The group therapy is shaped and configured by a number of therapeutic strategies which may include interpersonal, behavioral, cognitive-behavioral, psychodrama, and psycho-educational interventions.

The group is organized around an age group with some shared commonality of developmental, psychosocial, and sexual concerns. Groups are usually divided by age into early childhood, middle childhood, preadolescent, and adolescent groups. Group treatment with the younger sexually abusive child has been carried out with some success. Johnson and Berry (1989) have recommended both male and female co-therapists with groups of 5 or fewer children. They are generally grouped according to age i.e., 5 to 7, 8 to 10 and 11 to 13 years. The therapeutic intervention is structured and focused, emphasizing problem solving skills, social skills, anger management and cognitive strategies for controlling behavior. Thematic concerns include trust, betrayal, secrecy, guilt, labeling of affects, loss, helplessness, powerlessness, empowerment, sexual feelings, cognitive distortions, self-blame and self-esteem, etc. The families of these children are often seriously compromised.

An important goal in group therapy of the older juvenile offender is the achievement of a sense of group cohesion and peer acceptance. Adolescents are struggling to find their place in the peer group and to develop mutually reciprocal peer relations. They are concerned with intellectual and emotional self-expression, relationships with authority, sexual relations, moral choices, identificatory pathways, and the consolidation of a sense of self. Groups may be homogeneous or heterogeneous relative to categories of sexual offenses; open or closed; structured or unstructured. Initially, there is some advantage to the groups being structured around psycho-educational materials. The structured format provides a nidus around which the processing of emotionally arousing and threatening materials may come to the surface without being preoccupied with issues of trust (Schwartz, 1988).

The parents should receive concurrent parent counseling or be placed in a structured parent group with an emphasis on educational modules where upon they can discuss their own issues of victimization and more importantly focus on styles of interaction and management of their children’s sexual behavior.

Individual Psychotherapy

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

While historically individual therapy has been a valuable intervention, it has had limited value for the individual sex offender and probably should never be relied on as the only treatment model. The advantages of individual therapy is that it provides a greater sense of confidentiality and an opportunity to develop trust in the therapeutic process. Individual therapy does provide an opportunity to assist the individual in understanding his/her psychodynamics, unresolved developmental issues, conflicts, and resistance to therapeutic intervention. Most importantly individual therapy can be used to develop a therapeutic alliance which may be used to facilitate the offender's participation in other treatment modalities. There are several disadvantages to individual therapy. It is easier for the therapist to be manipulated. Denial is more easily sustained. There is less therapeutic confrontation. The sexual secrets are maintained within the therapeutic dyad. There is less opportunity to learn from others, i.e., victim empathy, offender characteristics, offense cycle, and interpersonal and social skills (Schwartz, 1988).

The therapy of juvenile sex offenders is generally characterized as one of firmness and confrontation alternating with a flexible and sympathetic stance (Muster, 1992). Confrontation is necessary to address the minimizations, denial, rationalization and the cognitive distortions which the offender presents to authority. This approach is balanced with a sensitive awareness of the offender's developmental, behavioral and emotional problems which not infrequently emanate from his/her own childhood history. Individual therapy is a valuable adjunct for those individuals who have been sexually abused. Frequently it is not until the offender becomes emotionally in contact with his/her own affects associated with his/her own history of sexual abuse and other maltreatment that they begin to extend empathy to their own victims. Individual therapy is often the treatment of choice for the younger sexually reactive abused child who has become sexually abusive. This is particularly true for children who manifest high levels of intrapsychic conflict, emotional distress, confusion and defensiveness around their own sexual victimization. Individual therapy allows for the development of a dyadic relationship in which issues of trust, sexually aggressive fantasies, shame and guilt can be worked out in the context of the therapeutic relationship. In most instances, the individual therapy will be juxtaposed to other interventions to possibly include family and group therapy, cognitive-behavioral therapy, and psychopharmacological interventions.

Family Therapy

It is within the family context that many of the offender's beliefs, myths, and cognitive distortions about sexuality, aggression, and gender evolve and have been maintained. Family

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

therapy provides an opportunity not only to understand the offender's development and coping strategies but provides an opportunity to correct cognitive distortions and family mythologies. Family therapy facilitates the learning of new ways of communicating and building a support system which will help interrupt the abuse cycle and ultimately be supportive to the offender's capacity for regulating and modulating sexual aggression (Schwartz, 1988, Sholevar and Schwoeri, 1999). Bischof et al. (1995) suggest that intervention strategies which have been proven effective in other delinquent groups may be effective with the families of adolescent sex offenders. Family therapy may be warranted in those instances where there is incest, especially when the sex offender remains in the family or will rejoin the nuclear family after treatment.

PSYCHOPHARMACOLOGICAL INTERVENTIONS

Selective Serotonin Reuptake Inhibitors (SSRI)

The SSRIs (Greenberg and Bradford, 1997) have been shown to diminish sexual drive, sexual arousal, and sexual preoccupations. Serotonergic dysfunctions has been associated with impulsivity, suicidal behavior and aggressive behaviors, (Brown et al., 1979; Coccaro et al., 1996; Linnoila et al., 1983). The SSRIs are effective in the treatment of obsessive-compulsive behaviors and their use has been recommended for those individuals with paraphiliac or nonparaphiliac compulsive sexual preoccupations. Nonparaphiliac compulsive sexual behavior (Coleman, 1992) is characterized by conventional and normative sexual behavior taken to a compulsive extreme. Others have referred to this condition as a "sexual addiction." Stein et al. (1992) in a retrospective study of the use of serotonin reuptake blockers in a small sample of "paraphiliac, nonparaphiliac sexual addictions and sexual obsessions" found those with sexual obsessions had the best response to medication. Clinical studies have suggested that fluvoxamine, clomipramine and buspirone may be effective (Greenberg and Bradford, 1997).

Fluoxetine has been the agent that has been most studied and there are a number of reports indicating that its use is associated with a reduction in paraphiliac behavior and non-paraphiliac sexual obsessions (Kafka and Prentky, 1992). Greenberg et al.(1996) carried out a retrospective study of 58 paraphiliacs treated over a 12 month period with either fluoxetine, fluvoxamine or sertraline and found paraphiliac fantasies to be markedly reduced. A 12 week controlled retrospective study of 95 paraphiliacs treated with SSRIs matched with a control group likewise found the severity and frequency of paraphiliac fantasies significantly reduced (Greenberg and Bradford, 1997).

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

Antiandrogens

There has been considerable interest in the use of antiandrogen drugs to decrease sexually aggressive behavior. The evidence for the role of testosterone in sexually aggressive behavior, however, is equivocal and conflicting (Archer, 1991; Bagatell et al., 1994; Brooks and Reddon, 1996). There are reports that higher levels of testosterone are associated with increased frequency of orgasm and sexual behaviors (Knussmann and Christiansen, 1986). Other studies have failed to demonstrate increasing levels of testosterone with increased overt or covert sexual behavior (Bagatell et al., 1994). The literature is confusing because of the confound between aggressive violence and sexuality. Brooks and Reddon (1996) studied 194 adolescents 15-17 years of age: 75 violent offenders, 102 non-violent offenders, and 17 sexual abusers. They found no differences between the non-violent and sexual abusers but significantly higher levels of testosterone in the violent group. Rada et al. (1976) found no differences in testosterone levels between rapists, child molesters and controls. When the rapists were divided in increasing gradations of violence the most violent rapists had the highest levels of testosterone.

Bradford and Bourget (1987) have reviewed the effects of antiandrogen drug treatment in sexual abusers and have observed that with decreased testosterone that there is a reduction in sexual drive, sexual desire, fantasy, and sexual intercourse 4 to 6 weeks after the onset of treatment. Studies have demonstrated the efficaciousness of medroxyprogesterone acetate (MPA), Depo-Provera, in adult sex offenders (Berlin, 1989; Berlin and Menicke, 1981). Depo-Provera substantially lowers serum testosterone and has been called a “sexual appetite suppressant” (Berlin, 1989).

While there is evidence that antiandrogen drug intervention decreases sexual preoccupation, sexual fantasy, and sexual behaviors, its use in adolescents has to be carefully considered (Prentky, 1997). Antiandrogen drugs such as cyproterone acetate (CPA) and MPA, may delay the onset of puberty. CPA has been associated with a number of undesirable side effects such as gynecomastia, hypersomnia, fatigue, depression and alterations in adrenal functioning while MPA has been related to increased weight gain, gastrointestinal upset, headaches, sleep disturbances, malaise and hyperglycemia (Katz, 1999). Luteinizing hormone-releasing hormone (LHRH) agonists have been shown in adults to reduce testosterone levels and concomitantly to significantly diminish deviant sexual fantasies and sexual behaviors (Katz, 1999; Rosler and Witztum, 1998).

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

These agents are reserved for the most severe sexual abusers and are generally discouraged for use in adolescents less than 17 years of age. Antiandrogen medications should never be used as an exclusive treatment for paraphiliac and aggressive sexual behaviors (Prentky, 1997).

There are considerable ethical and legal implications in the use of antiandrogen medications as they are not FDA approved for the treatment of sexual abusers. A signed informed consent statement, which is comprehensive and detailed in scope with full disclosure of the side effects, risks, and potential benefits, is necessary before initiating treatment.

AFTERCARE

A sexual abuser is never cured but is rehabilitated. There is a need to provide monitoring and follow-up with continuing services. After the termination of a course of therapeutic interventions, the offender is maintained in a spectrum of continuing services which resonate with the severity of his sexual misbehaviors and psychopathology and which may include community based outpatient treatment programs, specialized group homes, specialized foster care programs, and other specialized follow-up services.

CONFLICT OF INTEREST

As a matter of policy, some of the authors of these practice parameters are in active clinical practice and may have received income related to treatments discussed in these parameters. Some authors may be involved primarily in research or other academic endeavors and also may have received income related to treatments discussed in these parameters. To minimize the potential for these parameters to contain biased recommendations due to conflict of interest, the parameters were reviewed extensively by Work Group members, consultants, and Academy members; authors and reviewers were asked to base their recommendations on an objective evaluation of the available evidence; and authors and reviewers who believed that they might have a conflict of interest that would bias, or appear to bias, their work on these parameters were asked to notify the Academy.

SCIENTIFIC DATA AND CLINICAL CONSENSUS

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

Practice parameters are strategies for patient management, developed to assist clinicians in psychiatric decision-making. These parameters, based on evaluation of the scientific literature and relevant clinical consensus, describe generally accepted approaches to assess and treat specific disorders, or to perform specific medical procedures. The validity of scientific findings was judged by design, sample selection and size, inclusion of comparison groups, generalizability, and agreement with other studies. Clinical consensus was determined through extensive review by the members of the Work Group on Quality Issues, child and adolescent psychiatry consultants with expertise in the content area, the entire Academy membership, and the Academy Assembly Council.

These parameters are not intended to define the standard of care; nor should they be deemed inclusive of all proper methods of care or exclusive of other methods of care directed at obtaining the desired results. The ultimate judgment regarding the care of a particular patient must be made by the clinician in light of all the circumstances presented by the patient and his or her family, the diagnostic and treatment options available and available resources. Given inevitable changes in scientific information and technology, these parameters will be reviewed periodically and updated when appropriate.

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

REFERENCES

References marked with an asterisk are particularly recommended.

Abel G, Rouleau J, Cunningham-Rathner J (1986), Sexually Aggressive Behavior. In: *Forensic Psychiatry and Psychology: Perspectives and Standards for Interdisciplinary Practice*, Curran W, McGarry A, Shah S, eds. Philadelphia, PA: FA Davis, pp 289-313

Abel GG, Becker JV, Cunningham-Rathner J, Mittelman M, Rouleau JL (1988), Multiple paraphiliac diagnoses among sex offenders. *Bull Am Acad Psych Law* 16(2):153-168

Abel GG, Osborn C (1992), The paraphilias. The extent and nature of sexually deviant and criminal behavior. *Psychiatr Clin North Am* 15: 675-687

Ageton SS (1983), *Sexual Assault Among Adolescents*. Lexington, MA: Lexington Books
American Academy of Child and Adolescent Psychiatry (1997), Practice parameters for the psychiatric assessment of children and adolescents. *J Am Acad Child Adolesc Psychiatry* 36:4S-20S

American Psychiatric Association (1994), *DSM IV: Diagnostic and Statistical Manual of Mental Disorders, fourth edition*. Washington, DC: American Psychiatric Association

Anderson PB (1996), Correlates of college women's self-reports of heterosexual aggression. *Sexual Abuse: Journal of Research and Treatment* 8(2):121-131

Archer J (1991), The Influence of Testosterone on Human Aggression, *British J Psychol* 82:1-28

*Araji SK (1997), *Sexually Aggressive Children: coming to understand them*. Thousand Oaks, CA: Sage Publications

Awad GA, Levene J, Saunders E (1984), A clinical study of male adolescent sex offenders. *International Journal of Offender Therapy and Comparative Criminology* 28(2):105-116

Awad GA, Saunders EB (1989), Adolescent child molesters: Clinical observations. *Child Psychiatry Hum Dev* 19:195-206

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

Bagatell CJ, Heiman JR, Rivier JE, Bremner WJ (1994), Effects of Endogenous Testosterone and Estradiol on Sexual Behavior in Normal Young Men, *J Clin Endocrinol Metab* 78:711-716

Becker JV, Cunningham-Rathner J, Kaplan MS (1986), Adolescent Sexual Offenders: Demographics, criminal, and sexual histories, and recommendations for reducing future offenses. *J Interpersonal Violence* 1: 431-445

Becker JV, Kaplan MS, Cunningham-Rathner J, Kavoussi R (1986), Characteristics of adolescent incest sexual perpetrators: Preliminary findings. *Journal of Family Violence* 1:85-97

Becker JV (1988a), Adolescent Sex Offenders. *Behavior Therapist* 11:185-187

Becker JV, Hunter JA, Stein RM, Kaplan MS (1989), Factors associated with erection in adolescent sex offenders. *Journal of Psychopathology and Behavioral Assessment* 11:353-362

Becker JV (1990) Treating Adolescent Sexual offenders. *Professional Psychology: Research and Practice* 21: 362-365

Becker JV, Kaplan MS, Tenke CE, Tartaglino A (1991), The incidence of depressive symptomatology in juvenile sex offenders with a history of abuse. *Child Abuse Negl* 15:531-536

Becker JV, Kaplan MS, Tenke CE (1992), The relationship of abuse history, denial and erectile response profiles of adolescent sexual perpetrators. *Behavior Therapy* 23: 87-97

*Becker JV, Hunter JA (1993), Aggressive sex offenders. *Child and Adolescent Psychiatric Clinics of North America* 2:477-487

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

Becker JV, Harris CD, Sales BD (1993), Juveniles who commit sexual offenses: a critical review of research. In: *Sexual Aggression*, Nagayama-Hall GC, Hirschman R, Graham JR, Zaragoza T eds. Washington DC: Taylor and Francis, pp 215-228

*Becker JV (1994), Offenders: Characteristics and Treatment. *The Future of Children: Sexual Abuse of Children* 4:176-197

Becker JV, Hunter JA (1997), Understanding and Treating Child and Adolescent Sexual Offenders. In: *Advances in Clinical Child Psychology*, Ollendick TH, Prinz RJ, eds. New York: Plenum Press, pp 177-197

Berlin FS, Meinecke CF (1981), Treatment of sex offenders with antiandrogenic medication: Conceptualization, review of treatment modalities and preliminary findings. *Am J Psychiatry* 138:601-607

Berlin FS (1989), The Paraphilia and Depo-Provera: Some Medical, Ethical and Legal Considerations, *Bull Am Acad Psychiatry Law* 17: 233-239

Bischof GP, Stith SM, Whitney ML (1995), Family environments of adolescent sex offenders and other juvenile delinquents. *Adolescence* 30:157-170

Borduin CM, Henggeler SW, Blaske DM, Stein RJ (1990), Multisystemic treatment of adolescent sexual offenders. *International Journal of Offender Therapy and Comparative Criminology* 34:105-113

Bradford JM, Bourget D (1987), Sexually Aggressive Men. *Psychiatric Journal of the University of Ottawa* 12:169-175

Bremer JF (1992), Serious juvenile sex offenders: Treatment and long term follow-up. *Psychiatric Annals* 22:326-332

Brooks JH, Reddon JR (1996), Serum Testosterone in Violent and Nonviolent Young Offenders, *J Clin Psychol* 52:475-483

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

Brown GL, Goodwin FK, Ballenger JC, Goyer PF, Major LF (1979), Aggression in humans correlates with cerebrospinal fluid amine metabolites. *Psychiatric Research* 1:131-139

Carpenter DR, Peed SF, Eastman B (1995), Personality Characteristics of adolescent sexual offenders: A Pilot Study, *Sexual Abuse: Journal of Research and Treatment* 7(3) 195-203

Cautela JR (1966), Treatment of compulsive behavior by covert sensitization. *Psychological Record* 16:33-41

Coccaro EF, Kavoussi RJ, Sheline YI, Lisk JD, Csernansky JG (1996), Impulsive aggression in personality disorder correlates with tritiated paroxetine binding in the platelet. *Arch Gen Psychiatry* 53:531-536

Coleman E (1990), The Obsessive-Compulsive model for describing compulsive sexual behavior, *American Journal of Preventive Psychiatry and Neurology* 2(3):9-14

Coleman E (1992), Is your patient suffering from compulsive sexual behavior? *Psychiatric Annals* 22:320-325

Cooper CL, Murphy WD, Haynes MR (1996), Characteristics of Abused and Nonabused Adolescent Sexual Offenders. *Sexual Abuse: Journal of Research and Treatment* 8: 105-119

Dhawan S, Marshall WL (1996), Sexual Abuse Histories of Sexual Offenders. *Sexual Abuse: Journal of Research and Treatment* 8:7-15

Davis GE, Leitenberg H (1987), Adolescent Sex Offenders, *Psychol Bull* 101:417-427

Davis TC, Peck GQ, Storment JM (1993), Acquaintance Rape and the High School Student, *J Adolesc Health* 14:220-224

Deblinger E, McLeer SV, Atkins MS, Ralphe D, Foa E (1989), Post-traumatic stress in sexually abused, physically abused and nonabused children, *Child Abuse Negl* 13:403-408

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

- Doshay LJ (1943), *The Boy Sex Offender and His Later Career*. New York: Gruen and Stratton
- Dougher MJ (1988b), Clinical Assessment of the Sex Offenders. In: *A Practitioner's Guide to Treating the Incarcerated Male Sex Offender*, Schwartz BK, Cellini HR, eds. Washington DC: U.S. Department of Justice, National Institute of Corrections, US Government Printing Office, pp 77-84
- Dougher MJ (1988), Behavioral techniques to alter sexual arousal. In: *A Practitioner's Guide to Treating the Incarcerated Male Sex Offender*, Schwartz BK, Cellini HR, eds. Washington DC: US Department of Justice, National Institute of Corrections, US Government Printing Office, pp 109-113
- Dwyer SM (1997), Treatment Outcome Study: Seventeen Years after Sexual Offender Treatment, *Sexual Abuse: Journal of Research and Treatment* 9(2):149-160
- Elliott DS (1994a), *Youth violence: an overview*. Boulder CO: The Center for the Study and Prevention of Violence, University of Colorado Boulder Institute for Behavioral Sciences (F-693)
- Elliott DS (1994b), *The developmental course of sexual and non-sexual violence: results from a national longitudinal study*, Paper presented at the annual meeting of the American Association for the Treatment of Sexual Abusers, San Francisco CA
- Elliott DS (1994c), Serious violent offenders: Onset, developmental course, and termination. The American Society of Criminology 1993 Presidential Address. *Criminology* 32:1-21
- Equal Employment Opportunity Commission (1980), Title VII Guidelines on Sexual Harassment (Rules and Regulations, 74676-74-677). *Federal Register* 45:219
- Epps KJ, Haworth R, Swaffer T (1993), Attitudes toward woman and rape among male adolescents convicted of sexual versus nonsexual crimes. *J Psychol* 127: 501-506

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

Farrell JK, O'Brien R (1988), *Sexual Offenses by Youth in Michigan: Data Implications and Policy Recommendations. Report to the Michigan Legislature*. Detroit, MI: Safer Society Resources of Michigan

*Fehrenbach PA, Smith W, Monastersky C, Deisher RW (1986), Adolescent sexual offenders: offender and offense characteristics. *Am J Orthopsychiatry* 56 (suppl 2):225-233

Fehrenbach P, Monastersky C (1988), Characteristics of female adolescent sex offenders, *Am J Orthopsychiatry* 58:148-151

Ford ME, Linney JA (1995), Comparative analysis of juvenile sexual offenders, violent nonsexual offenders and status offenders. *Journal of Interpersonal Violence* 10:56-70

Freund K (1963), A laboratory method for diagnosing predominance of homo or hetero erotic interest in the male. *Behav Res Ther* 1:85-93

*Friedrich WN, Grambsch P, Broughton D, Kuiper J, Beilke RL (1991), Normative sexual behavior in children. *Pediatrics* 88: 456-464

Friedrich WN (1993), Sexual Victimization and Sexual Behavior in Children: A Review of Recent Literature *Child Abuse Negl* 17:59-66

Friedrich WN (1995), *Psychotherapy with Sexually Abused Boys: an integrated approach*. Thousand Oaks, CA: Sage Publications

Friedrich WN (1997), *CSBI: Child Sexual Behavior Inventory. Professional Manual*. Odessa, Florida Psychological Assessment Resources, Inc.

Friedrich WN, Grambsch P, Damon L, Hewitt S, Koverla C, Lang R et al. (1992), The child sexual behavior inventory: Normative and clinical contrasts. *Psychological Assessment* 4: 303-311

Furby L, Weinrott MR, Blackshaw L (1989), Sex offender recidivism: A review. *Psychol Bull* 105: 3-30

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

- Gerdes KE, Gourley MM, Cash MC, (1995), Assessing juvenile sex offenders to determine adequate levels of supervision, *Child Abuse Negl*, 19:953-961
- Gil E, Johnson TC (1993), *Sexualized Children: Assessment and Treatment of Sexualized Children and Children Who Molest*. Rockville, MD: Launch Press
- Gilby R, Wolf L, Goldberg B (1989), Mentally retarded adolescent sex offenders, a survey and pilot study, *Canadian Journal of Psychiatry* 34:542-548
- Gray AS, Pithers WD (1993), Relapse prevention with sexually aggressive adolescents and children: Expanding treatment and supervision. In: *The Juvenile Sex Offender*, Barbaree HE, Marshall WL, Hudson SM, eds. New York: Guilford Press, pp 289-319
- Gray SR (1995), A comparison of verbal satiation and minimal arousal conditioning to reduce deviant arousal in the laboratory. *Sexual Abuse: A Journal of Research and Treatment* 7(2): 143-154
- Gray A, Busconi A, Houchens P, Pithers WD (1997), Children with sexual behavior problems and their caregivers: Demographics, functioning and clinical patterns. *Sexual Abuse: Journal of Research and Treatment* 9(4) 267-290
- Green R (1988), Psycho-Educational Modules. In: *A Practitioner's Guide to Treating the Incarcerated Male Sex Offender*. Schwartz BK, Cellini HR, eds. Washington, DC: US Department of Justice, National Institute of Corrections, US Government Printing Office, pp 95-100
- Green A (1999), Female Sexual Offenders. In: *Sexual Aggression*, Shaw JA, ed. Washington DC: American Psychiatric Press
- Green AH, Kaplan MS (1994), Psychiatric impairment and childhood victimization experiences in female child molesters. *J Am Acad Child Adolesc Psychiatry* 33:954-961

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

- *Greenberg DM, Bradford JMW, Curry S, O'Rourke A (1996), A comparison of treatment of paraphilias with three serotonin reuptake inhibitors: A Retrospective study, *Bulletin of the American Academy of Psychiatry and the Law* 24: 525-532
- *Greenberg DM, Bradford JMW (1997), Treatment of the Paraphiliac Disorders: A Review of the Role of the Selective Serotonin Reuptake Inhibitors, *Sexual Abuse: Journal of Research and Treatment* 9(4):349-360
- Groth NA, Hobson WF, Lucey KP, Pierre JS (1981), Juvenile Sex Offenders: Guidelines for Treatment, *International Journal Offender Therapy and Comparative Criminology*, 25:65-72
- *Hall GC (1995), Sexual offender recidivism revisited: A meta-analysis of recent treatment studies. *J Consult Clin Psychol* 63: 802-809
- Hampton HL (1995), Care of the woman who has been raped. *N Engl J Med* 332:234-237
- Harvey MR, Herman JL (1992), The Trauma of Sexual Victimization: Feminist Contributions to Theory, Research, and Practice. *PTSD Research Quarterly* 3(3): 1-7
- Hastings T, Anderson SJ, Hemphill P (1997), Comparisons of daily stress, coping, problem behavior and cognitive distortions in adolescent sex offenders and conduct disordered youth, *Sexual Abuse: Journal of Research and Treatment* 9(1): 29-42
- Herkov MJ, Gynther MD, Thomas S, Myers WC, (1996), MMPI differences among adolescent inpatient, rapists, sodomists and sexual abusers, *J Pers Assess* 66(1):81-90
- Holmes WC, Slap GB (1998), Sexual abuse of boys: definition, prevalence, correlates, sequelae, and management. *JAMA* 280:1855-1865
- Hunter JA (1991), A comparison of the psychosocial maladjustment of adult males and females sexually molested as children. *Journal of Interpersonal Violence* 6:205-217

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

Hunter JA, Becker JV, Kaplan M, Goodwin DW (1991), Reliability and discriminative utility of the Adolescent Cognitions Scale for juvenile sexual offenders. *Annals of Sex Research* 4:281-286

Hunter JA, Becker JV, Kaplan MS (1995), The Adolescent Sexual Interest Card Sort: Test-Retest Reliability and Concurrent Validity in Relation to Phallometric Assessment. *Archives of Sexual Behavior* 24:555-561

Hunter JA, Lexier LJ (1998), Ethical and Legal Issues in the Assessment and Treatment of Juvenile Sex Offenders. *Child Maltreatment* 3:339-348

Hunter JA, Becker JV (1999), Motivators of adolescent sex offenders and treatment perspectives. In: *Sexual Aggression*, Shaw JA, ed. Washington DC: American Psychiatric Press, pp 211-235

Hunter JA, Figueredo AJ (1999), Factors associated with Treatment Compliance in a Population of Juvenile Sexual Offenders. *Sexual Abuse: A Journal of Research and Treatment* 11:49-67

Jacobs WL, Kennedy WA, Meyer JB (1997), Juvenile Delinquents: A between-group comparison study of sexual and nonsexual offenders. *Sexual Abuse: Journal of Research and Treatment* 9:201-217

Johnson TC (1988), Child perpetrators-children who molest other children: preliminary findings. *Child Abuse Negl* 12:219-229

Johnson TC (1989), Female child perpetrators: children who molest other children. *Child Abuse Neglect* 13: 571-585

Johnson TC, Berry C (1989), Children who molest: a treatment program. *Journal of Interpersonal Violence* 4(2):185-203

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

- Johnson T (1999), The development of sexual behavior problems in childhood. In: *Sexual Aggression*, Shaw JA, ed. Washington, DC: American Psychiatric Press
- Johnston L, Ward T (1996), Social Cognition and Sexual Offending: A Theoretical Framework, *Sexual Abuse: Journal of Research and Treatment* 8(1):55-80
- Kaemingk KL, Koselka M, Becker JV, Kaplan MS (1995), Age and Adolescent Sexual Offender Arousal. *Sexual Abuse: Journal of Research and Treatment* 7(4):249-257
- Kafka MP, Prentky R (1992), Fluoxetine treatment of non-paraphiliac sexual addictions and paraphilias in men, *Journal of Clinical Psychiatry* 53: 351-358
- Kahn TJ, Chambers HJ (1991), *Assessing Reoffense Risk with Juvenile Sex Offenders*. *Child Welfare* 70:333-345
- Kaplan MS, Green A (1995), Incarcerated female sexual offenders: A comparison of sexual histories with eleven female nonsexual offenders. *Sexual Abuse: Journal of Research and Treatment* 7(4):287-300
- *Katz D (1999), Psychopharmacological interventions with adolescent and adult sex offenders. In: *Sexual Aggression*, Shaw JA, ed. Washington DC: American Psychiatric Press, pp 305-326
- Katz RC (1990), Psychosocial adjustment in adolescent child molesters. *Child Abuse Negl* 14:567-575
- Kavoussi RJ, Kaplan M, Becker JV (1988), Psychiatric diagnoses in adolescent sex offenders. *J Am Acad Child Adolesc Psychiatry* 27: 241-243
- *Kendall-Tackett KA, Williams LM, Finkelhor D (1993), Impact of sexual abuse on children: A review and synthesis of recent empirical studies. *Psychol Bull* 113: 164-180

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

- Knight RA, Prentky RA (1993), Exploring characteristics for classifying juvenile sex offenders. In: *The Juvenile Sex Offender*, Barbaree HE, Marshall WL, Hudson SM, eds. New York: Guilford Press, pp 45-83
- Knopp FH (1991), *The Youthful Sex Offender: The Rationale and Goals of Early Intervention and Treatment*. Orwell VT: Safer Society Program and Press
- Knopp FH, Longo RF, Stevenson WF (1992), *Nationwide Survey of Juvenile and Adult Sex-Offender Treatment Program and Models*. Orwell, VT: Safer Society Program and Press
- Knopp FH, Lackey LB (1987), *Female Sexual Abusers: A Summary of Data from 44 Treatment Providers*. Orwell, VT: Safer Society Program and Press
- Knussmann R, Christiansen K, Couwenbergs C (1986), Relations between sex hormone levels and sexual behavior in men. *Arch Sex Behav* 15:429-445
- Kobayashi J, Sales BD, Becker JV, Figueredo AJ, Kaplan M, (1995), Perceived Parental deviance, parent child bonding, child abuse and child sexual aggression, *Sexual Abuse: Journal of Research and Treatment* 7(1):25-44
- Levin SM, Stava L (1987), Personality Characteristics of Sex Offenders: A Review, *Arch Sex Behav* 16(1):57-79
- Lewis DO, Shankok SS, Pincus JH (1979), Juvenile male sexual assaulters. *Am J Psychiatry* 136(suppl 9):1194-1196
- Linnoila M, Virkkunen M, Scheinin M, Nuutila A, Rimon R, Goodwin FK (1983), Low cerebrospinal fluid 5-hydroxyindolacetic acid concentration differentiates impulsive from non-impulsive violent behavior. *Life Sci* 33: 2609-2614
- Longo RE (1982), Sexual Learning and Experience among adolescent sexual offenders, *International Journal of Offender Therapy and Comparative Criminology* 26(3): 235-241

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

Maletzky BM (1974), "Assisted" covert sensitization in the treatment of exhibitionism, *Journal of Consulting and Clinical Psychology* 42: 34-40

Maletzky BM (1997), Editors Note. *Sexual Abuse: A Journal of Research and Treatment* 9:147

Maletzky BM (1996), Denial of treatment or treatment of denial. *Sexual Abuse: A Journal of Research and Treatment* 8: 1-5

Marshall WL, Barbaree HE (1988), An outpatient treatment program for child molesters. *Ann NY Acad Sci* 528: 205-214

Mathews R, Matthews JK, Speltz K (1989), *Female sex offenders: An exploratory study*.
Brandon, VT: Safer Society Press

*Mathews R, Hunter JA, Vuz J (1997), Juvenile female sexual offenders: Clinical characteristics and treatment issues. *Sexual Abuse: A Journal of Research and Treatment* 9:187-199

McClellan J, McCurry C, Ronnei M, Adams J, Eisner A, Storck M (1996), Age of Onset of Sexual Abuse: Relationship to Sexually Inappropriate Behaviors. *J Am Acad Child Adolesc Psychiatry* 35:1375-1383

McClellan J, McCurry C, Ronnei M, Adams J, Storck M, Eisner A., Smith C, (1997), Relationship Between Sexual Abuse, Gender, and Sexually Inappropriate Behaviors in Seriously Mentally Ill Youths. *J Am Acad Child Adolesc Psychiatry* 36:959-965

McConaghy N, Armstrong MS, Blaszczyński MA, Kidson W (1989), Resistance to treatment of adolescent sex offenders. *Arch Sex Behav* 18:97-107

McCurry C, McClellan J, Adams J, Ronnei M, Storck M, Eisner A, Breiger D (1998), Sexual Behavior Associated with Low Verbal IQ in Youth who have Severe Mental Illness. *Mental Retardation* 36: 23-30

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

- Miner MH (1997), How can we conduct treatment outcome research, *Sexual Abuse: A Journal of Research and Treatment* 9(2):95-110
- Money J (1986), *Lovemaps: Clinical concepts of sexual/erotic health and pathology, paraphilia, and gender transposition in childhood, adolescence, and maturity*. New York: Irvington Publishers
- Murphy WD, Coleman EM, Haynes MR (1983), Treatment and evaluation issues with the mentally retarded sex offender. In: *The Sexual Aggressor: Current Perspectives on Treatment*, Greer JG, Stuart IR eds. New York: Van Nostrand Reinhold
- Muster NJ (1992), Treating the adolescent victim-turned-offender. *Adolescence* 27:441-450
- National Committee to Prevent Child Abuse (1998), *Current Trends in Child Abuse Reporting and Fatalities: The Results of the 1994 Annual Fifty State Survey*. Chicago: NCPCA (Available from NCPCA, 332 S. Michigan Avenue, Suite 1600, Chicago, IL, 60604)
- National Task Force on Juvenile Sexual Offending (1993), National Adolescent Perpetrator Network Revised Report. *Juvenile and Family Court Journal* 44(suppl 4): 1-121
- Nichols HR, Molinder I (1984), *Manual for the Multiphasic Sex Inventory*. Tacoma WA: Crime and Victim Psychology Specialists
- Office of Juvenile Justice and Delinquency Prevention (1994), *Juveniles and Violence*. Washington, DC: US Department of Justice, Office of Justice Programs, Office of Juvenile Justice and Delinquency Prevention (Fact Sheet #19)
- Office of Juvenile Justice and Delinquency Prevention (1994a), *Violent Families and Youth Violence*. Washington, DC: US Department of Justice, Office of Justice Programs, Office of Juvenile Justice and Delinquency Prevention (Fact Sheet #21)
- Packard WS, Rosner R (1985), Psychiatric evaluations of sexual offenders. *J Forensic Sci* 30:715-720

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

Pithers WD, Gibat CC, Marlatt GA, Marques JK (1983), Relapse prevention: a self-control model of treatment and maintenance of change for sexual aggression. In: *The Sexual Aggressor: Current Perspective on Treatment*, Greer JG, Stuart IR, eds. New York: Van Nostrand Reinhold, pp 214-239

Pithers WD, Beal L, Cumming G, Turner R, Young W (1988a), Relapse prevention. In: *A Practitioner's Guide to Treating the Incarcerated Male Sex Offender*, Schwartz BK, Cellini HR, eds. Washington DC: US Department of Justice, National Institute of Corrections, US Government Printing Office, pp 123-140

Pithers WD, Beal LS, Buell MM, Cummings GF, Kashima KM (1988b), Relapse prevention of sexual aggression. In: *Human Sexual Aggression: Current Perspectives*, Prentky RA, Quinsey VL, eds. New York: New York Academy of Sciences, pp 244-260

*Pithers WD, Gray AS (1996), Utility of relapse prevention in treatment of sexual abusers. *Sexual Abuse: Journal of Research and Treatment* 8(3):223-230

Ponton LE (1996), Sexual harassment of children and adolescent girls. In: *Sexual Harassment in the Workplace and Academia: Psychiatric Issues*, Shrier D, ed. Washington DC: American Psychiatric Press, Inc., pp. 181-201

Postman N (1994), *The Disappearance of Childhood*. New York: Vintage Books

Prentky R, Burgess AW (1990), Rehabilitation of child molesters: a cost-benefit analysis. *Am J Orthopsychiatry* 60:108-117

Prentky, RA (1997), Arousal reduction in sexual offenders: a review of antiandrogen interventions. *Sexual Abuse: Journal of Research and Treatment* 9(4): 335-347

Prentky RA, Knight RA, Lee AFS (1997), *Research Report: Child sexual molestation: research issues*. Washington, DC: U.S. Government Printing Office

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

- Proulx J, Lusignan R, McKibben A (1996), Relationship between affective components and sexual behaviors in sexual aggressors. *Sexual Abuse: A Journal of Research and Treatment* 8(4):279-289
- Rada RT, Laws DR, Kellner R (1976), Plasma testosterone in the rapist. *Psychosom Med* 38:257-268
- Rasmussen LA (1999), Factors related to recidivism among juvenile sexual offenders. *Sexual Abuse: A Journal of Research and Treatment* 11(1):69-85
- Ross J, Loss P (1991), Assessment of the juvenile sex offender. In: *Juvenile Sex Offending*, Ryan GD, Lane SL, eds. Lexington MA: Lexington Books, pp 199-252
- Rosenfeld AA, Wasserman S (1993), Sexual development in the early school-aged child. *Child and Adolescent Psychiatric Clinics of North America* 2:393-406
- Rosler A, Witztum, E (1998), Treatment of men with paraphilia with a long-acting analogue of gonadotropin releasing hormone. *New Engl J Med* 338:416-422
- Rubinstein M, Yeager C, Goodstein C, Lewis DO (1993), Sexually assaultive male juveniles: a follow-up. *Am J Psychiatry* 150:262-265
- Ryan G, Lane S, Davis J, Isaac C, (1987), Juvenile sex offenders: development and correction. *Child Abuse Negl* 11:385-395
- Ryan G, Miyoshi TJ, Metzner JL, Krugman RD, Fryer GE, (1996), Trends in a national sample of sexually abusive youths. *J Am Acad Child Adolesc Psychiatry* 35(Suppl 1):17-25
- Saunders E, Awad GA, White G (1986), Male adolescent sexual offenders: the offender and the offense. *Can J Psychiatry* 31:542-549
- Saunders EB, Awad GA (1988), Assessment, management, and treatment planning for male adolescent sexual offenders. *Am J Orthopsychiatry* 58:571-579

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

- Shoor M, Speed MH, Bartelt C (1966), Syndrome of the adolescent child molester. *Am J Psychiatry* 122: 783-789
- Schlank AM, Shaw T (1996), Treating sexual offenders who deny their guilt: a pilot study. *Sexual Abuse: Journal of Research and Treatment* 8(1):17-23
- Schwartz B (1988), Interpersonal techniques in treating sex offenders. In: *A Practitioner's Guide to Treating the Incarcerated Male Sex Offender*, Schwartz BK, Cellini HR, eds. Washington DC: US Department of Justice, National Institute of Corrections, US Government Printing Office, pp 101-107
- Schwartz BK (1992), Effective treatment techniques for sex offenders. *Psychiatric Annals* 22:315-319
- Sefarbi R (1990), Admitters and deniers among adolescent sex offenders and their families: a preliminary study. *Am J Orthopsychiatry* 60: 460-465
- Segal ZV, Stermac L (1984), A measure of rapist's attitudes towards women. *Inter Jour Law Psychiatry* 7:437-440
- Shaw J, Campo-Bowen AE, Applegate B, Perez D, Antoine LB, Hart EL, Lahey BB, Testa RJ, Devaney A (1993), Young boys who commit serious sexual offenses: demographics, psychometrics, and phenomenology. *Bull Am Acad Psychiatry Law* 21:399-408
- Shaw JA, Applegate B, Rothe E (1996), Psychopathology and personality disorders in adolescent sex offenders. *American Journal of Forensic Psychiatry* 17(4):19-37
- Shaw, JA (1999), Sexually aggressive behavior. In: *Sexual Aggression*, Shaw JA, ed. Washington DC: American Psychiatric Press, pp 3-40
- Shaw T, Funderburk JR, Schlank AM (1999), Cognitive-behavioral treatment strategies and the integrated model for adolescent perpetrators of sexual abuse. In: *Sexual Aggression*, Shaw JA, ed. Washington, DC: American Psychiatric Press, pp 275-304

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

- Sholevar GP, Schwoeri LD (1999), Sexual aggression within the family. In: *Sexual Aggression*, Shaw JA, ed. Washington, DC: American Psychiatric Press, pp 75-106
- Siegel LJ, Senna JJ (1988), *Juvenile Delinquency*, Third Edition. San Francisco: West Publishing Company
- Smets AC, Cebula CM (1987), A group treatment program for adolescent sex offenders: five steps toward resolution. *Child Abuse Negl* 11: 247-254
- Smith WR, Monastersky C (1986), Assessing juvenile sexual offender's risk for reoffending. *Criminal Justice and Behavior* 13(2): 115-140
- Stein DJ, Hollander E, Anthony DT, Schneier FR, Fallon BA, Liebowitz MR, Klein DF, (1992), Serotonergic medications for sexual obsessions, sexual addictions, and paraphilias. *J Clin Psychiatry* 53:267-271
- Stermac LE, Sheridan P (1993), The developmentally disabled adolescent sex offender. In: *The Juvenile Sex Offender*, Barbaree HE, Marshall WL, Hudson SM, eds. New York: Guilford Press pp. 235-242
- Straus S (1988), Sexual harassment in the school: legal implications for principals. *National Association of Secondary Schools Principals Bulletin*, pp 93-97, March 1988 (quoted in Ponton LE (1995), Sexual harassment of children and adolescent girls. In: *Sexual Harassment in the Workplace and Academia: Psychiatric Issues*, Schrier D, ed. Washington, DC: American Psychiatric Press, pp 107-128)
- Tartar RE, Alterman AI, Hegedus AM, Katz-Garris (1983), Cognitive capacities of juvenile, non-violent and sexual offenders. *J Nerv Ment Dis* 171:364-367
- Truscott D (1993), Adolescent offenders: comparison for sexual, violent, and property offenses. *Psychol Rep* 73:657-658

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

Watkins B, Bentovim A (1992), The sexual abuse of male children and adolescents: a review of current research. *J Child Psychol Psychiatry* 33:197-248

White JW, Koss MP (1993), Adolescent sexual aggression within heterosexual relationships: prevalence, characteristics and causes. In: *The Juvenile Sex Offender*, Barbaree HE, Marshall WL, Hudson SM, eds. New York: Guilford Press, pp 182-202

Widom CS, (1989), The cycle of violence. *Science* 244:160-166

Worling JR (1995), Adolescent sibling-incest offenders: differences in family and individual functioning when compared to adolescent non sibling sex offenders. *Child Abuse Negl* 19:633-643

Yates A (1982), Children eroticized by incest. *Am J Psychiatry* 139:482-485

Zgourides G, Monto M, Harris R (1994), Prevalence of prior adult sexual contact in a sample of adolescent male sex offenders. *Psychol Rep* 75:1042

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

TABLE 1

DEFINITIONS

Sexually Abusive Behavior *	Sexual behavior which occurs without consent; without equality or as a result of coercion
Sexual Offense *	Sexually violating/exploiting behavior breaching societal norms and moral codes resulting in physical or psychological harm; a violation of federal, state or municipal law, statute or ordinance
Rape *	To seize or take by force for sexual gratification
Sodomy *	Usually indicates anal intercourse but may include oral as well
Sexual Harassment **	Unwelcomed sexual attention which may consist of sexual overtures, requests, advances and verbal or physical conduct of a sexual nature.
Sexual offender *	An individual who has committed an act of sexual aggression breaching societal norms and moral codes, violated federal, state, municipal law, statute or ordinance

AMERICAN ACADEMY OF CHILD AND ADOLESCENT PSYCHIATRY

<p>Paraphilia ***</p>	<p>Recurrent, intense sexually arousing fantasies, sexual urges, or behaviors generally involving 1) nonhuman objects, 2) the suffering or humiliation of oneself or one's partner or 3) children or other nonconsenting persons that occurs over at least a six month period</p>
<p>Pedophilia ***</p>	<p>Sexual activities focused on a prepubescent child generally younger than 13 years of age</p>
<p>Sexually Reactive Children ****</p>	<p>Children who display sexually inappropriate behavior in response to sexual abuse or exposure to explicit sexual stimuli</p>

*National Task Force on Juvenile Sexual Offending, 1993

** Equal Employment Opportunity Commission, Title VII, 1980

***APA, 1994; In order for a youth to be diagnosed as a Pedophile he must be sixteen years or older and at least five years older than the child.

**** Yates, 1982; Gil and Johnson, 1993.

Table 2

**ASSESSMENT OF THE SEXUAL ABUSIVE INCIDENT:
WHAT ONE WANTS TO LEARN FROM THE CLINICAL INTERVIEW**

- Degree of cooperation.
- Honesty and forthrightness of the abuser.
- Degree of acceptance of responsibility for his sexual offenses.
- Degree of remorse and regret.
- Relationship between the abuser and the victim.
- Age difference between the abuser and the victim.
- Characteristics of the sexual aggressive behavior.
- Frequency and duration of the sexual aggressive behavior.
- Precipitating factors that led to the sexual offense.
- Premeditated or impulsive.
- Characteristics of the victim that attracted the offender.
- Nature and extent of the coercive behaviors.
- Behaviors prior to, during and post sexual offense.
- Affect states prior to, during and post sexual offenses.
- Verbal interchange with the victim.
- Attempts to avoid detection.
- Understanding of the effects of his sexual behavior on the victim.
- Insight into the wrongfulness of his sexual behavior.
- Understanding of the consequences of his behavior.